

A Fractional Analogue of Brooks' Theorem

Andrew D. King ^{*} Linyuan Lu [†] Xing Peng [‡]

January 15, 2013

Abstract

Let $\Delta(G)$ be the maximum degree of a graph G . Brooks' theorem states that the only connected graphs with chromatic number $\chi(G) = \Delta(G) + 1$ are complete graphs and odd cycles. We prove a fractional analogue of Brooks' theorem in this paper. Namely, we classify all connected graphs G such that the fractional chromatic number $\chi_f(G)$ is at least $\Delta(G)$. These graphs are complete graphs, odd cycles, C_8^2 , $C_5 \boxtimes K_2$, and graphs whose clique number $\omega(G)$ equals the maximum degree $\Delta(G)$. Among the two sporadic graphs, the graph C_8^2 is the square graph of cycle C_8 while the other graph $C_5 \boxtimes K_2$ is the strong product of C_5 and K_2 . In fact, we prove a stronger result; if a connected graph G with $\Delta(G) \geq 4$ is not one of the graphs listed above, then we have $\chi_f(G) \leq \Delta(G) - \frac{2}{67}$.

1 Introduction

The chromatic number of graphs with bounded degrees has been studied for many years. Brooks' theorem perhaps is one of the most fundamental results; it is included by many textbooks on graph theory. Given a simple connected graph G , let $\Delta(G)$ be the maximum degree, $\omega(G)$ be the clique number, and $\chi(G)$ be the chromatic number. Brooks' theorem states that $\chi(G) \leq \Delta(G)$ unless G is a complete graph or an odd cycle. Reed [10] proved that $\chi(G) \leq \Delta(G) - 1$ if $\omega(G) \leq \Delta(G) - 1$ and $\Delta(G) \geq \Delta_0$ for some large constant Δ_0 . This excellent result was proved by probabilistic methods, and Δ_0 is at least hundreds. Before this result, Borodin and Kostochka [1] made the following conjecture.

Conjecture [1]: Suppose that G is a connected graph. If $\omega(G) \leq \Delta(G) - 1$ and $\Delta(G) \geq 9$, then we have

$$\chi(G) \leq \Delta(G) - 1.$$

If the conjecture is true, then it is best possible since there is a K_8 -free graph $G = C_5 \boxtimes K_3$ (actually K_7 -free, see Figure 1) with $\Delta(G) = 8$ and $\chi(G) = 8$.

Here we use the following notation of the strong product. Given two graphs G and H , the *strong product* $G \boxtimes H$ is the graph with vertex set $V(G) \times V(H)$, and (a, x) is connected to (b, y) if one of the following holds

- $a = b$ and $xy \in E(H)$,

^{*}Simon Fraser University, Burnaby, British Columbia, Canada (andrew.d.king@gmail.com). This author was supported in part by an EBCO/Ebbich Postdoctoral Scholarship and the NSERC Discovery Grants of Pavol Hell and Bojan Mohar.

[†]University of South Carolina, Columbia, SC 29208, (lu@math.sc.edu). This author was supported in part by NSF grant DMS 1000475.

[‡]University of South Carolina, Columbia, SC 29208, (pengx@mailbox.sc.edu). This author was supported in part by NSF grant DMS 1000475.




Figure 1: The graph $C_5 \square K_3$.

- $ab \in E(G)$ and $x = y$,
- $ab \in E(G)$ and $xy \in E(H)$.

Reed's result [10] settled Borodin and Kostochka's conjecture for sufficiently large $\Delta(G)$, but the cases with small $\Delta(G)$ are hard to cover using the probabilistic method.

In this paper we consider a fractional analogue of this problem. The fractional chromatic number $\chi_f(G)$ can be defined as follows. A *b-fold coloring* of G assigns a set of b colors to each vertex such that any two adjacent vertices receive disjoint sets of colors. We say a graph G is *a:b-colorable* if there is a b -fold coloring of G in which each color is drawn from a palette of a colors. We refer to such a coloring as an *a:b-coloring*. The *b-fold coloring number*, denoted by $\chi_b(G)$, is the smallest integer a such that G has an *a:b-coloring*. Note that $\chi_1(G) = \chi(G)$. It was shown that $\chi_{a+b}(G) \leq \chi_a(G) + \chi_b(G)$. The *fractional chromatic number* $\chi_f(G)$ is $\lim_{b \rightarrow \infty} \frac{\chi_b(G)}{b}$.

By the definition, we have $\chi_f(G) \leq \chi(G)$. The fractional chromatic number can be viewed as a relaxation of the chromatic number. Many problems involving the chromatic number can be asked again using the fractional chromatic number. The fractional analogue often has a simpler solution than the original problem. For example, the famous $\omega - \Delta - \chi$ conjecture of Reed [9] states that for any simple graph G , we have

$$\chi(G) \leq \left\lceil \frac{\omega(G) + \Delta(G) + 1}{2} \right\rceil.$$

The fractional analogue of $\omega - \Delta - \chi$ conjecture was proved by Molloy and Reed [8]; they actually proved a stronger result with ceiling removed, i.e.,

$$\chi_f(G) \leq \frac{\omega(G) + \Delta(G) + 1}{2}. \quad (1)$$

In this paper, we classify all connected graphs G with $\chi_f(G) \geq \Delta(G)$.

Theorem 1 *A connected graph G satisfies $\chi_f(G) \geq \Delta(G)$ if and only if G is one of the following*

1. a complete graph,
2. an odd cycle,

3. a graph with $\omega(G) = \Delta(G)$,

4. C_8^2 ,

5. $C_5 \boxtimes K_2$.

For the complete graph K_n , we have $\chi_f(K_n) = n$ and $\Delta(K_n) = n - 1$. For the odd cycle C_{2k+1} , we have $\chi_f(C_{2k+1}) = 2 + \frac{1}{k}$ and $\Delta(C_{2k+1}) = 2$. If G is neither a complete graph nor an odd cycle but contains a clique of size $\Delta(G)$, then we have

$$\Delta(G) \leq \omega(G) \leq \chi_f(G) \leq \chi(G) \leq \Delta(G). \quad (2)$$

The last inequality is from Brooks' theorem. The sequence of inequalities above implies $\chi_f(G) = \Delta(G)$.

If G is a vertex-transitive graph, then we have [11]

$$\chi_f(G) = \frac{|V(G)|}{\alpha(G)},$$

where $\alpha(G)$ is the independence number of G . Note that both graphs C_8^2 and $C_5 \boxtimes K_2$ are vertex-transitive and have the independence number 2. Thus we have

$$\chi_f(C_8^2) = 4 = \Delta(C_8^2) \quad \text{and} \quad \chi_f(C_5 \boxtimes K_2) = 5 = \Delta(C_5 \boxtimes K_2).$$



Figure 2: The graph C_8^2 and $C_5 \boxtimes K_2$.

Actually, Theorem 1 is a corollary of the following stronger result.

Theorem 2 *Assume that a connected graph G is neither C_8^2 nor $C_5 \boxtimes K_2$. If $\Delta(G) \geq 4$ and $\omega(G) \leq \Delta(G) - 1$, then we have*

$$\chi_f(G) \leq \Delta(G) - \frac{2}{67}.$$

Remark: In the case $\Delta(G) = 3$, Heckman and Thomas [5] conjectured that $\chi_f(G) \leq 14/5$ if G is triangle-free. Hatami and Zhu [4] proved $\chi_f(G) \leq 3 - \frac{3}{64}$ for any triangle-free graph G with $\Delta(G) \leq 3$. The second and third authors showed an improved result $\chi_f(G) \leq 3 - \frac{3}{43}$ in the previous paper [7]. Thus we need only consider the cases $\Delta(G) \geq 4$. For any connected graph G with sufficiently large $\Delta(G)$ and $\omega(G) \leq \Delta(G) - 1$, Reed's result [10] $\chi(G) \leq \Delta(G) - 1$ implies $\chi_f(G) \leq \Delta(G) - 1$. The method introduced in [4] and strengthened in [7], has a strong influence on this paper. The readers are encouraged to read these two papers [4, 7].

Let $f(k) = \inf_G \{\Delta(G) - \chi_f(G)\}$, where the infimum is taken over all connected graphs G with $\Delta(G) = k$ and not one of the graphs listed in Theorem 1. Since $\chi_f(G) \geq \omega(G)$, by taking a graph with $\omega(G) = \Delta(G) - 1$, we have $f(k) \leq 1$. Theorem 2 says $f(k) \geq \frac{2}{67}$ for any $k \geq 4$. Reed's result [10] implies $f(k) = 1$ for sufficiently large k . Heckman and Thomas [5] conjectured $f(3) = 1/5$. It is an interesting problem to determine the value of $f(k)$ for small k . Here we conjecture $f(4) = f(5) = \frac{1}{3}$. If Borodin and Kostochka's conjecture is true, then $f(k) = 1$ for $k \geq 9$.

Theorem 2 is proved by induction on k . Because the proof is quite long, we split the proof into the following two lemmas.

Lemma 1 *We have $f(4) \geq \frac{2}{67}$.*

Lemma 2 *For each $k \geq 6$, we have $f(k) \geq \min\{f(k-1), \frac{1}{2}\}$. We also have $f(5) \geq \min\{f(4), \frac{1}{3}\}$.*

It is easy to see the combination of Lemma 1 and Lemma 2 implies Theorem 2. The idea of reduction comes from the first author, who pointed out $f(k) \geq \min\{f(k-1), \frac{1}{2}\}$ for $k \geq 7$ based on his recent results [6]. The second and third authors originally proved $f(k) \geq \frac{C}{k^3}$ (for some $C > 0$) using different method in the first version; they also prove the reductions at $k = 5, 6$, which are much harder than the case $k \geq 7$. We do not know whether a similar reduction exists for $k = 4$.

The rest of this paper is organized as follows. In section 2, we will introduce some notation and prove Lemma 2. In section 3 and section 4, we will prove $f(4) \geq \frac{2}{67}$.

2 Proof of Lemma 2

In this paper, we use the following notation. Let G be a simple graph with vertex set $V(G)$ and edge set $E(G)$. The *neighborhood* of a vertex v in G , denoted by $\Gamma_G(v)$, is the set $\{u : uv \in E(G)\}$. The *degree* $d_G(v)$ of v is the value of $|\Gamma_G(v)|$. The *independent set* (or *stable set*) is a set S such that no edge with both ends in S . The *independence number* $\alpha(G)$ is the largest size of S among all the independent sets S in G . When $T \subset V(G)$, we use $\alpha_G(T)$ to denote the independence number of the induced subgraph of G on T . Let $\Delta(G)$ be the maximum degree of G . For any two vertex-sets S and T , we define $E_G(S, T)$ as $\{uv \in E(G) : u \in S \text{ and } v \in T\}$. Whenever G is clear under context, we will drop the subscript G for simplicity.

If S is a subset of vertices in G , then *contracting* S means replacing vertices in S by a single fat vertex, denoted by $\underline{S}$, whose incident edges are all edges that were incident to at least one vertex in S , except edges with both ends in S . The new graph obtained by contracting S is denoted by G/S . This operation is also known as *identifying vertices of* S in the literature. For completeness, we allow S to be a single vertex or even the empty set. If S only consists of a single vertex, then $G/S = G$; if $S = \emptyset$, then G/S is the union of G and an isolated vertex. When S consists of 2 or 3 vertices, for convenience, we write G/uv for $G/\{u, v\}$ and G/uvw for $G/\{u, v, w\}$; the fat vertex will be denoted by $\underline{uv}$ and $\underline{uvw}$, respectively. Given two disjoint subsets S_1 and S_2 , we can contract S_1 and S_2 sequentially. The order of contractions does not matter; let $G/S_1/S_2$ be the resulted graph. We use $G - S$ to denote the subgraph of G induced by $V(G) - S$.

In order to prove Lemma 2, we need use the following theorems due to King [6].

Theorem 3 (King [6]) *If a graph G satisfies $\omega(G) > \frac{2}{3}(\Delta(G) + 1)$, then G contains a stable set S meeting every maximum clique.*

Theorem 4 (King [6]) For a positive integer k , let G be a graph with vertices partitioned into cliques $V_1, \dots, V_r$. If for every i and every $v \in V_i$, v has at most $\min\{k, |V_i| - k\}$ neighbors outside V_i , then G contains a stable set of size r .

Lemma 3 Suppose that G is a connected graph with $\Delta(G) \leq 6$ and $\omega(G) \leq 5$. Then there exists an independent set meeting all induced copies of K_5 and $C_5 \boxtimes K_2$.

Proof: We first show that there exists an independent set meeting all copies of K_5 . If G contains no K_5 , then this is trivial. Otherwise, we can apply Theorem 3 to get the desired independent set since $\omega(G) > \frac{2}{3}(\Delta(G) + 1)$ is satisfied.

Now we prove the Lemma by contradiction. Suppose the Lemma is false. Let G be a minimum counterexample (with the smallest number of vertices). For any independent set I , let $C(I)$ be the number of induced copies of $C_5 \boxtimes K_2$ in $G - I$. Among all independent sets which meet all copies of K_5 , there exists one such independent set I such that $C(I)$ is minimized.

Since $C(I) > 0$, there is an induced copy of $C_5 \boxtimes K_2$ in $G - I$; we use H to denote it. In $C_5 \boxtimes K_2$, there is a unique perfect matching such that identifying the two ends of each edge in this matching results a C_5 . An edge in this unique matching is called a *canonical* edge. We define a new graph G' as follows: First we contract all canonical edges in H to get a C_5 , where its vertices are called *fat* vertices. Second we add five edges turning the C_5 into a K_5 . Observe that each vertex in this C_5 can have at most two neighbors in $G - H$ and $\Delta(G') \leq 6$. We will consider the following four cases.

Case 1: There is a K_6 in the new graph G' . Since the original graph G is K_6 -free, the K_6 is formed by the following two possible ways.

Subcase 1a: This K_6 contains 5 fat vertices. By the symmetry of H , there is an induced C_5 in H such that the vertices in C_5 contain a common neighbor vertex v in $G \setminus V(H)$, see Figure 3. Since H is K_5 -free, we can find x, y in this C_5 such that x, y is a non-edge. Let $I' := (I \setminus \{v\}) \cup \{x, y\}$; I' is also an independent set. Observe that v is not in any K_5 in $G - I'$. Thus the set I' is also an independent set and meets every K_5 in G . Since $C_5 \boxtimes K_2$ is a 5-regular graph, any copy of $C_5 \boxtimes K_2$ containing v must contain at least one of x and y . Thus, $C(I') < C(I)$. Contradiction!




Figure 3: Subcase 1a.




Figure 4: Subcase 1b.

Subcase 1b: This K_6 contains 4 fat vertices. Let u, v be the other two vertices. By the symmetry of H , there is a unique way to connect u and v to H as shown by Figure 4. Since uv is an edge, one of u and v is not in I . We assume $u \notin I$. Let $\{x, y\} \subset \Gamma_G(v) \cap V(H)$ as shown in Figure 4 and $I' = I \setminus \{v\} \cup \{x, y\}$. Observe that I' is an independent set and v is not in a K_5 in $G - I'$. Thus I' is an independent set meeting each K_5 in G . Since each $C_5 \boxtimes K_2$ containing v must contain one of x and y . Thus $C(I') < C(I)$. Contradiction!

Case 2: There is a K_5 intersecting H with 4 vertices. Let v be the vertex of this K_5 but not in H , see Figure 5. We have two subcases.

Subcase 2a: The vertex v has another neighbor y in H but not in this K_5 . Since H is K_5 -free, we can select a vertex x in this K_5 such that xy is not an edge of G . Let $I' := I \setminus \{v\} \cup \{x, y\}$. Note that v is not in a K_5 in $G - I'$, and I' is an independent set. Thus I' is an independent set meeting each K_5 in G . Since any $C_5 \boxtimes K_2$ containing v must contain one of x and y , we have $C(I') < C(I)$. Contradiction!

Subcase 2b: All neighbors of v in H are in this K_5 . Let x be any vertex in this K_5 other than v , and $I' := I \setminus \{v\} \cup \{x\}$. In this case, there is only one K_5 containing v . Thus, I' is also an independent set meeting every copy of K_5 in G . Observe that $\Gamma_G(v) \setminus \{x\}$ is disconnected. If $v \in H' = C_5 \boxtimes K_2$, then $\Gamma_G(v) \cap H'$ is connected. Thus v is not in a $C_5 \boxtimes K_2$ in $G - I'$ and $C(I') < C(I)$. Contradiction!




Figure 5: Case 2.




Figure 6: Case 3.

Case 3: There is an induced subgraph H' isomorphic to $C_5 \boxtimes K_2$ such that H' and H are intersecting, see Figure 6. Since $V(H) \cap V(H') \neq \emptyset$ and $H \neq H'$, we can find a canonical edge uv of H and a canonical edge uv' of H' such that $v \notin V(H')$ and $v' \notin V(H)$. If vv' is a non-edge, then let $I' := I \setminus \{v'\} \cup \{u\}$. It is easy to check I' is still an independent set. We also observe that any possible K_5 containing v' must also contain u . Thus, I' meets every copy of K_5 in G . We have v' in no $C_5 \boxtimes K_2$ in $G - I'$ since vv' is not an edge. We therefore get $C(I') < C(I)$. Contradiction! If vv' is an edge, then locally there are two K_5 intersecting at u, v , and v' ; say the other four vertices are x_1, x_2, y_1, y_2 , where two cliques are $\{x_1, x_2, u, v, v'\}$ and $\{y_1, y_2, u, v, v'\}$, see Figure 6. Let $I' = I \cup \{x_1, y_1\} \setminus \{v'\}$. Note that I' is an independent set and v' is not in a K_5 in $G - I'$. Thus I' is an independent set meeting each K_5 in G . Observe that any copies of $C_5 \boxtimes K_2$ containing v' must contain one of x_1 and y_1 ; we have $C(I') < C(I)$. Contradiction!

Case 4: This is the remaining case, G' is K_6 -free. We have $\omega(G') \leq 5$ and $|V(G')| < |V(G)|$. By the minimality of G , there is an independent set I' of G' meeting every copy of K_5 and $C_5 \boxtimes K_2$. In I' , there is a unique vertex x of the K_5 obtained from contracting canonical edges of H . Let uv be the canonical edge corresponding to x . Let $I'' = I' \setminus \{x\} \cup \{u\}$, we get an independent set I'' of G . Note that any $v \in H \setminus \{u\}$ is not in any K_5 of $G - I''$ by Case 2 as well as not in any $C_5 \boxtimes K_2$ of $G - I''$ by Case 3. Thus I'' hits each K_5 in G and $C(I'') = 0$. Contradiction! $\square$

The following lemma extends Theorem 3 when $\omega(G) = 4$; a similar result was proved independently in [2].

Lemma 4 *Let G be a connected graph with $\Delta(G) \leq 5$ and $\omega(G) \leq 4$. If $G \neq C_{2l+1} \boxtimes K_2$ for some $l \geq 2$, then there is an independent set I hitting all copies of K_4 in G .*

Proof: We will prove it by contradiction. If the lemma is false, then let G be a minimum counterexample. If G is K_4 -free, then there is nothing to prove. Otherwise, we consider the clique graph $\mathcal{C}(G)$, whose edge set is the set of all edges appearing in some copy of K_4 . Because of $\Delta(G) = 5$, here are all possible connected component of $\mathcal{C}(G)$.

1. $C_t \boxtimes K_2$ for $t \geq 4$. If this type occurs, then every vertex in $C_t \boxtimes K_2$ has degree 5; thus, this is the entire graph G . If t is even, then we can find an independent set I meeting every K_4 . If t is odd, then it is impossible to find such an independent set. However, this graph is excluded from the assumption of the Lemma.
2. $P_t \boxtimes K_2$ for $t \geq 3$. In this case, all internal vertices have degree 5 while the four end vertices have degree 4. Consider a new graph G' which is obtained by deleting all internal vertices and adding four edges to make the four end vertices as a K_4 . It is easy to check $\Delta(G') \leq 5$ and $\omega(G') \leq 4$. Since $|G'| < |G|$, there is an independent set I of G' meeting every copy of K_4 in G' . Note that there is exactly one end vertex in I . Observe that any one end vertex can be extended into a maximal independent set meeting every copy of K_4 in $P_t \boxtimes K_2$. Thus, we can extend I to an independent set I' of G such that I' meets every copy of K_4 in G . Hence, this type of component does not occur in $\mathcal{C}(G)$.
3. There are four other types listed in Figure 7.




Figure 7: All types of components in the clique graph $\mathcal{C}(G)$.

For each component C_i in $\mathcal{C}(G)$, let V_i be the set of common vertices in all K_4 's of C_i ; for the leftmost figure in Figure 7, V_i is the set of all 4 vertices; for the middle two figures, V_i is the set of bottom three vertices; for the rightmost figure, V_i consists of the left-bottom vertex and the middle-bottom vertex. Note that all V_i 's are pairwise disjoint. Let G' be the induce subgraph of G on $\cup_i V_i$. Note that G' does not contains any vertex in $C_i \setminus V_i$. By checking each type, we find out that for each i and each $v \in V_i$, v has at most $\min\{2, |V_i| - 2\}$ neighbors outside V_i in G' (not in $G!$). Applying Theorem 4 to G' , we conclude that there exists an independent set I of G' meeting every V_i ; thus I meets every K_4 in G . Contradiction!

□

Lemma 5 *Let G be a connected graph with $\Delta(G) \leq 5$ and $\omega(G) \leq 4$. If $G \neq C_{2l+1} \boxtimes K_2$ for some $l \geq 2$, then there exists an independent set meeting all induced copies of K_4 and C_8^2 .*

Proof: We will use proof by contradiction. Suppose the Lemma is false. Let G be a minimum counterexample (with the smallest number of vertices). For any independent set I , let $C(I)$ be the number of induced copies of C_8^2 in $G - I$. Among all independent sets which meet all copies of K_4 , there exists an independent set I such that $C(I)$ is minimized. Since $C(I) > 0$, let H be a copy of C_8^2 in $G - I$. The vertices of H are listed by u_i for $i \in \mathbb{Z}_8$ anticlockwise such that $u_i u_j$ is an edge of H if and only if $|i - j| \leq 2$. The vertex u_{i+4} is the *antipode* of u_i for any $i \in \mathbb{Z}_8$.

Case 1: There exists a vertex $v \notin V(H)$ such that v has five neighbors in H . By the Pigeonhole Principle, $\Gamma(v)$ contains a pair of antipodes. Without loss of generality, say $u_0, u_4 \in \Gamma(v)$. If the other three neighbors of v do not form a triangle, then we let $I' := I \setminus \{v\} \cup \{u_0, u_4\}$; note that v is not in any K_4 of $G - I'$. Thus I' is an independent set meeting every copy of K_4 . Since every copy of C_8^2 containing v must contain one of u_0 and

u_4 , we have $C(I') < C(I)$. Contradiction! Hence, the other three neighbors of v must form a triangle. Without loss of generality, we can assume that the three neighbors are u_1 , u_2 , and u_3 . Now we let $I'' := I \setminus \{v\} \cup \{u_0, u_3\}$; note that $v \notin K_4 \subset G - I'$. Thus I'' is also an independent set meeting every copy of K_4 of G . Since every copy of C_8^2 containing v must contain one of u_0 and u_3 , we have $C(I'') < C(I)$. Contradiction!

Case 2: There exists a vertex $v \notin V(H)$ such that v has exactly four neighbors in H . Since H is K_4 -free, we can find $u_i, u_j \in \Gamma(v) \cap V(H)$ such that $u_i u_j$ is a non-edge. Let $I' := I \setminus \{v\} \cup \{u_i, u_j\}$; I' is also an independent set. Note that $\Gamma(v) \setminus \{u_i, u_j\}$ can not be a triangle, v is not in any $K_4 \subset G - I'$. Thus I' meets every copy of K_4 . Since every copy of C_8^2 containing v must contain one of u_i and u_j , we have $C(I') < C(I)$. Contradiction!

Case 3: There exists a vertex $v \notin V(H)$ such that v has exactly three neighbors in H . If the 3 neighbors do not form a triangle, then choose $u_i, u_j \in \Gamma(v) \cap V(H)$ such that $u_i u_j$ is a non-edge. Note that $\Gamma(v) \setminus \{u_i, u_j\}$ can not be a triangle; v is not in any $K_4 \subset G - I'$. Let $I' := I \setminus \{v\} \cup \{u_i, u_j\}$; I' is also an independent set meeting every copy of K_4 . Since every copy of C_8^2 containing v must contain one of u_i and u_j , we have $C(I') < C(I)$. Contradiction! Else, the three neighbors form a triangle; let u_i be one of them and $I' := I \setminus \{v\} \cup \{u_i\}$; v is not in any $K_4 \subset G - I'$. Thus I' is an independent set meeting every copy of K_4 . Note that $\Gamma(v) \setminus \{u_i\}$ has only two vertices in H . The induced graph on $\Gamma(v) \setminus \{u_i\}$ is disconnected. However, for any vertex v in $H' = C_8^2$, the subgraph induced by $\Gamma_G(v) \cap H'$ is a P_4 . There is no C_8^2 in $G - I'$ containing v . Thus, $C(I') < C(I)$. Contradiction!

Case 4: Every vertex outside H can have at most 2 neighbors in H . We identify each pair of antipodes of H to get a new graph G' from G . After identifying, H is turned into a K_4 ; where the vertices of this K_4 are referred as fat vertices.

Subcase 4a: $G' \neq C_{2l+1} \boxtimes K_2$. Observe $\Delta(G') \leq 5$. We claim G' is K_5 -free. Suppose not. Since every vertex in H has at most one neighbor outside H , then each fat vertex can have at most two neighbors outside H . Recall that the original graph G is K_5 -free. If G' has some K_5 , then this K_5 contains either 3 or 4 fat vertices. Let w be one of the other vertices in this K_5 . We get w has at least three neighbors in H . However, this is covered by Case 1, Case 2, or Case 3. Thus, G' is K_5 -free. Since $|G'| < |G|$, by the minimality of G , G' has an independent set I' meeting every copy of K_4 and C_8^2 in G' . There is exactly one fat vertex in I' . Now replacing this fat vertex by its corresponding pair of antipodal vertices, we get an independent set I'' ; we assume the pair of antipodal vertices are u_2 and u_6 . It is easy to check that I'' is an independent set of G . Next we claim any $v \in V(H) \setminus \{u_2, u_6\}$ is neither in a $K_4 \subset V(G) - I''$ nor in a $C_8^2 \subset V(G) - I''$. Suppose there is some v such that $v \in K_4 \subset G - I''$. Recall each $v \in V(H)$ has at most one neighbor outside H and H is K_4 -free; there is some $w \notin V(H)$ such that w has at least three neighbors in H . This is already considered by Case 1, Case 2, or Case 3. We are left to show that $v \notin C_8^2 \subset G - I''$ for each $v \in V(H) \setminus \{u_2, u_6\}$. If not, there exists a copy H' of C_8^2 in $G - I''$ containing v . Note H' is 4-regular, any vertex in H' can have at most one neighbor in I'' ; in particular, $v \neq u_0, u_4$. Without loss of generality, we assume $v = u_3$. Then there is a vertex $w \notin V(H)$ such that $u_3 w$ is an edge, see Figure 8. Observe that the neighborhood of each vertex of an induced C_8^2 is is a P_4 . Since $u_1 u_4$ and $u_1 u_5$ are two non-edges, we have $w u_1$ being an edge. Observe $\Gamma_G(u_1) = \{u_7, u_0, u_2, u_3, w\}$. Since $u_2 \notin H'$, we have $u_0 \in H'$; u_0 has two neighbors (u_2 and u_6) outside H' , contradiction! Therefore, I'' meets every copy of K_4 and C_8^2 in G . Contradiction!

Subcase 4b: $G' = C_{2l+1} \boxtimes K_2$. The graph G can be recovered from G' . It consists of an induced subgraph $H = C_8^2$ and an induced subgraph $P_{2l-1} \boxtimes K_2$. For each vertex u in H , there is exactly one edge connecting it to one of the four end vertices of $P_{2l-1} \boxtimes K_2$; for each end vertex v of $P_{2l-1} \boxtimes K_2$, there are exactly two edges connecting v to the vertices in H . First, we take any maximum independent set I' of $P_{2l-1} \boxtimes K_2$. Observe that I' has exactly two end points of $P_{2l-1} \boxtimes K_2$; so I' has exactly four neighbors in H . In the remaining four




Figure 8: Subcase 4a.

vertices of H , there exists a non-edge $u_i u_j$ since H is K_4 -free. Let $I := I' \cup \{u_i, u_j\}$. Clearly I is an independent set of G meeting every copy of K_4 and C_8^2 . Contradiction! $\square$

We are ready to prove Lemma 2.

Proof of Lemma 2: We need prove for $k \geq 5$ and any connected graph G with $\Delta(G) = k$ and $\omega(G) \leq k - 1$ satisfies

$$\chi_f(G) \leq k - \min \left\{ f(k-1), \frac{1}{2} \right\}. \quad (3)$$

If $\omega(G) \leq k - 2$, then by inequality (1), we have

$$\chi_f(G) \leq \frac{\Delta(G) + \omega(G) + 1}{2} \leq k - \frac{1}{2}.$$

Thus, inequality (3) is satisfied. From now on, we assume $\omega(G) = \Delta(G) - 1$.

For $\Delta(G) = k \geq 6$ and $\omega(G) = k - 1$, the condition $\omega(G) > \frac{2}{3}(\Delta(G) + 1)$ is satisfied. By Theorem 3, G contains an independent set meeting every maximum clique. Extend this independent set to a maximal independent set and denote it by I . Note that $\Delta(G - I) \leq k - 1$ and $\omega(G - I) \leq k - 2$.

Case 1: $k \geq 7$. From the definition of $f(k - 1)$, we have $\chi_f(G - I) \leq \Delta(G - I) - f(k - 1)$. Thus,

$$\chi_f(G) \leq \chi_f(G - I) + 1 \leq k - 1 - f(k - 1) + 1 = k - f(k - 1).$$

Thus, we have $f(k) \geq \min\{f(k - 1), 1/2\}$.

Case 2: $k = 6$. By Lemma 3, we can find an independent set meets every copy of K_5 and $C_5 \boxtimes K_2$; we extend this independent set as a maximal independent set I . Note that $G - I$ contains no induced subgraph isomorphic $C_5 \boxtimes K_2$. We have $\chi_f(G - I) \leq 5 - f(5)$; it implies $\chi_f(G) \leq 6 - f(5)$. Thus, $f(6) \geq \min\{f(5), 1/2\}$ and we are done.

Case 3: $k = 5$. If $G = C_{2l+1} \boxtimes K_2$ for some $l \geq 3$; then G is vertex-transitive and $\alpha(G) = l$. It implies that

$$\chi_f(G) = \frac{|V(G)|}{\alpha(G)} = 4 + \frac{2}{l} \leq 5 - \frac{1}{3}.$$

If $G \neq C_{2l+1} \boxtimes K_2$, then by Lemma 5, we can find an independent set meeting every copy of K_4 and C_8^2 ; we extend it as a maximal independent set I . Note that $G - I$ contains no induced subgraph isomorphic C_8^2 . We have $\chi_f(G - I) \leq 4 - f(4)$; it implies $\chi_f(G) \leq 5 - f(4)$. Thus, $f(3) \geq \min\{f(4), 1/3\}$ and we are finished. $\square$

3 The case $\Delta(G) = 4$

To prove $f(4) \geq \frac{2}{67}$, we will use an approach which is similar to those in [4, 7]. We will construct 133 4-colorable auxiliary graphs, and from these colorings we will construct a 134-fold coloring of G using 532 colors.

It suffices to prove that the minimum counterexample does not exist.

Let G be a graph with the smallest number of vertices and satisfying

1. $\Delta(G) = 4$ and $\omega(G) \leq 3$;
2. $\chi_f(G) > 4 - \frac{2}{67}$;
3. $G \neq C_8^2$.

By the minimality of G , each vertex in G has degree either 4 or 3. To prove Lemma 1, we will show $\chi_f(G) \leq 4 - \frac{2}{67}$, which gives us the desired contradiction.

For a given vertex x in $V(G)$, it is easy to color its neighborhood $\Gamma_G(x)$ using 2 colors. If $d_G(x) = 3$, then we pick a non-edge S from $\Gamma_G(x)$ and color the two vertices in S using color 1. If $d_G(x) = 4$ and $\alpha(\Gamma_G(x)) \geq 3$, then we pick an independent set S in $\Gamma_G(x)$ of size 3 and assign the color 1 to each vertex in S . If $d_G(x) = 4$ and $\alpha(\Gamma_G(x)) = 2$, then we pick two disjoint non-edges S_1 and S_2 from $\Gamma_G(x)$; we assign color 1 to each vertex in S_1 and color 2 to each vertex in S_2 .

The following Lemma shows that G has a key property, which eventually implies that this local coloring scheme works simultaneously for x in a large subset of $V(G)$.

Lemma 6 *For each $x \in V(G)$ with $d_G(x) = 4$ and $\alpha(\Gamma_G(x)) = 2$, there exist two vertex-disjoint non-edges $S_1(x), S_2(x) \subset \Gamma_G(x)$ satisfying the following property. If we contract $S_1(x)$ and $S_2(x)$, then the resulting graph $G/S_1(x)/S_2(x)$ contains neither K_5^- nor G_0 . Here K_5^- is the graph obtained from K_5 by removing one edge and G_0 is the graph shown in Figure 9.*




Figure 9: The graph G_0 .

The proof of this lemma is quite long and we will present its proof in section 4.

For each vertex x in G , we associate a small set of vertices $S(x)$ selected from $\Gamma_G(x)$ as follows. If $d_G(x) = 3$, then let $S(x)$ be the endpoints of a non-edge in $\Gamma_G(x)$ and label the vertices in $S(x)$ as 1; if $d_G(x) = 4$ and $\alpha(\Gamma_G(x)) \geq 3$, then let $S(x)$ be any independent set of size 3 in $\Gamma_G(x)$ and label all vertices in $S(x)$ as 1; if $d_G(x) = 4$ and $\alpha(\Gamma_G(x)) = 2$, then

let $S(x) = S_1(x) \cup S_2(x)$, where $S_1(x)$ and $S_2(x)$ are guaranteed by Lemma 6; we label the vertices in $S_1(x)$ as 1 and the vertices in $S_2(x)$ as 2. For any $x \in V(G)$, we have $|S(x)| = 2, 3$, or 4.

The following definitions depend on the choice of $S(*)$, which is assumed to be fixed through this section. For $v \in G$ and $j \in \{1, 2, 3\}$, we define

$$N_G^j(v) = \{u \mid \text{there is a path } vv_0 \dots v_{j-2}u \text{ in } G \text{ of length } j \text{ such that } v_0 \in S(v) \text{ and } v_{j-2} \in S(u)\}.$$

We now define $N_G^j(u)$ for $j \in \{4, 5, 7\}$; each $N_G^j(u)$ is a subset of the j th neighborhood of u . For $j = 4$, $v \in N_G^4(u)$ if $d_G(u) = 4$, $\alpha(\Gamma_G(u)) = 2$, u and v are connected as shown in Figure 10; otherwise $N_G^4(u) = \emptyset$. In Figure 10, w is connected to one of the two vertices in $S_2(u)$. Similarly, in Figure 11 and 12, a vertex is connected to a group of vertices means it is connected to any vertex in this group.

For $j = 5$, $v \in N_G^5(u)$ if $d_G(w) = 4$, $\alpha(\Gamma_G(w)) = 2$ for $w \in \{u, v\}$ and u and v are connected as shown in Figure 11; otherwise $N_G^5(u) = \emptyset$.

For $j = 7$, $v \in N_G^7(u)$ if $d_G(w) = 4$, $\alpha(\Gamma_G(w)) = 2$ for $w \in \{u, v\}$ and u and v are connected as shown in Figure 12; otherwise $N_G^7(u) = \emptyset$.




Figure 10: 4-th neighborhood. Figure 11: 5-th neighborhood. Figure 12: 7-th neighborhood.

Note that for $j \in \{1, 2, 3, 5, 7\}$, $v \in N_G^j(u)$ if and only if $u \in N_G^j(v)$; but this does not hold for $j = 4$. We have the following lemma.

Lemma 7 For $u \in V(G)$ such that $d_G(u) = 4$ and $\alpha(\Gamma_G(u)) = 2$, we have $|N_G^1(u) \cup N_G^2(u) \cup N_G^3(u) \cup N_G^4(u) \cup N_G^5(u) \cup N_G^7(u)| \leq 96$.

Proof: It is clear that $|N_G^1(u) \cup N_G^2(u) \cup N_G^3(u)| \leq 4 + 8 + 8 \times 3 = 36$. We next estimate $|N_G^4(u)|$. In Figure 10, observe that w is connected to one vertex of $S_2(u)$ and $w \notin \Gamma_G(u)$. For a fixed u , there are at most four choices for w , at most three choices for z , and at most three choices for v . Therefore, we have $|N_G^4(u)| \leq 4 \times 3 \times 3 = 36$.

Let us estimate $|N_G^5(u)|$. In Figure 11, for a fixed u , we have four choices for w and two choices for z . Fix a z . Assume $\Gamma_G(z) \setminus \{w\} = \{a, b, c\}$. Let $T_1 = \{a, b\}$, $T_2 = \{b, c\}$, and $T_3 = \{a, c\}$. We have the following claim.

Claim There are at most three $v \in N_G^5(u)$ such that for each v we have $\Gamma_G(z) \cap \Gamma_G(v) = T_i$ for some $1 \leq i \leq 3$ as shown in Figure 11.

Proof of the claim: For each $1 \leq i \leq 3$, there are at most three $v \in N_G^5(u)$ such that $\Gamma_G(z) \cap \Gamma_G(v) = T_i$ as shown in Figure 11 since each vertex in T_i has at most three neighbors other than z . If the claim is false, then there is $1 \leq i \neq j \leq 3$ such that $\Gamma_G(z) \cap \Gamma_G(v_i) = T_i$ and $\Gamma_G(z) \cap \Gamma_G(v'_i) = T_i$ for some $v_i, v'_i \in N_G^5(u)$, and $\Gamma_G(z) \cap \Gamma_G(v_j) = T_j$ for some $v_j \in N_G^5(u)$, where v_i, v'_i, v_j are distinct. Without loss of generality, we assume $\Gamma_G(z) \cap \Gamma_G(v_1) = \Gamma_G(z) \cap \Gamma_G(v'_1) = T_1$ for $v_1, v'_1 \in N_G^5(u)$, and $\Gamma_G(z) \cap \Gamma_G(v_2) = T_2$ for some $v_2 \in N_G^5(u)$, see Figure 13. Observe that $\Gamma_G(b) = \{v_1, v'_1, v_2, z\}$. Since $\Gamma_G(z) \cap \Gamma_G(v_1) = T_1$

as shown in Figure 11, a and one of b 's neighbors form $S_i(v_1)$ for some $i \in \{1, 2\}$; we assume it is $S_1(v_1)$. Note $\{z, v_1, v'_1\} \subset \Gamma_G(a)$. Thus $S_1(v_1) = \{a, v_2\}$ and $v_2 \in \Gamma_G(v_1)$. Similarly, we can show $S_1(v'_1) = \{a, v_2\}$ and $v_2 \in \Gamma_G(v'_1)$. Now, observe that $\Gamma_G(v_2) = \{v_1, v'_1, b, c\}$. Since $\Gamma_G(z) \cap \Gamma_G(v_2) = T_2$ as shown in Figure 11, b and one of neighbors of v_2 form $S_i(v_2)$ for some $i \in \{1, 2\}$; we assume $i = 1$. Because $\{v_1, v'_1\} \subset \Gamma_G(b)$, then $S_1(v_2) = \{b, c\}$. However, b and c are not in the same independent set in the definition of $N_G^5(u)$, see Figure 11. This is a contradiction and this case can not happen. The claim follows.




Figure 13: The picture for the claim.

Therefore, $|N_G^5(u)| \leq 4 \times 2 \times 3 = 24$.

In Figure 12, for a fixed u , we have two choices for the edge e , one choice for w , two choices for z , and three choices for the edge f . Fix a z . By considering the degrees of the endpoints of f , there is at most one f and at most one $v \in N_G^7(u)$ such that $|\Gamma_G(f) \cap \Gamma_G(v)| = 4$ as shown in Figure 12. Therefore, we have $|N_G^7(u)| \leq 2 \times 2 \times 1 = 4$.

Last, we estimate $|N_G^5(u) \cup N_G^7(u)|$. If there is some $v \in N_G^7(u)$, then we observe that there are at most five z 's (see Figure 11). We get the number of $v \in N_G^5(u)$ is at most $5 \times 3 = 15$. In this case, we have

$$|N_G^5(u) \cup N_G^7(u)| \leq 4 + 15 < 24.$$

If $N_G^7(u) = \emptyset$, then also we have

$$|N_G^5(u) \cup N_G^7(u)| \leq 24.$$

Therefore

$$|N_G^1(u) \cup N_G^2(u) \cup N_G^3(u) \cup N_G^4(u) \cup N_G^5(u) \cup N_G^7(u)| \leq 36 + 36 + 24 = 96.$$

□

Based on the graph G , we define an auxiliary graph G^* on vertex set $V(G)$. The edge set is defined as follows: $uv \in E(G^*)$ precisely if either $u \in N_G^1(v) \cup N_G^2(v) \cup N_G^3(v) \cup N_G^4(v) \cup N_G^5(v) \cup N_G^7(v)$, or $v \in N_G^4(u)$. We have the following lemma.

Lemma 8 *The graph G^* is 133-colorable.*

Proof: Let σ be an increasing order of $V(G^*)$ satisfying the following conditions.

- 1: For u and v such that $d_G(u) = 3$ and $d_G(v) = 4$, we have $\sigma(u) < \sigma(v)$.
- 2: For u and v such that $d_G(u) = d_G(v) = 4$, $\alpha(\Gamma_G(u)) \geq 3$, and $\alpha(\Gamma_G(v)) = 2$, we have $\sigma(u) < \sigma(v)$.

We will color $V(G^*)$ according to the order σ . For each v , we have the following estimate on the number of colors forbidden to use for v .

- 1: For v such that $d_G(v) = 3$, the number of colors forbidden to use for v is at most $|N_G^1(v) \cup N_G^2(v) \cup N_G^3(v)| \leq 3 + 9 + 27 = 39$.
- 2: For v such that $d_G(v) = 4$ and $\alpha(\Gamma_G(v)) \geq 3$, the number of colors forbidden to use for v is at most $|N_G^1(v) \cup N_G^2(v) \cup N_G^3(v)| \leq 3 + 9 + 27 = 39$.
- 3: For v such that $d_G(v) = 4$ and $\alpha(\Gamma_G(v)) = 2$, the number of colors forbidden to use for v is at most $|N_G^1(v) \cup N_G^2(v) \cup N_G^3(v) \cup N_G^4(v) \cup N_G^5(v) \cup N_G^7(v)| + |N_G^4(v)| \leq 96 + 36 = 132$ by Lemma 7.

Therefore, the greedy algorithm shows G^* is 133-colorable. $\square$

Let X be a color class of G^* . We define a new graph $G(X)$ by the following process.

1. For each $x \in X$, if $|S(x)| = 2$ or $|S(x)| = 3$, then we contract $S(x)$ as a single vertex, delete the vertices in $\Gamma_G(v) \setminus S(v)$, and keep label 1 on the new vertex; if $|S(x)| = 4$, i.e., $S(x) = S_1(x) \cup S_2(x)$, then we contract $S_1(x)$ and $S_2(x)$ as single vertices and keep their labels. After that, we delete X . Let H be the resulting graph.
2. Note that $\Gamma_H(x) \cap \Gamma_H(y) = \emptyset$ and there is no edge from $\Gamma_H(x)$ to $\Gamma_H(y)$ for any $x, y \in X$ as X is a color class.
3. We identify all vertices with label i as a single vertex w_i for $i \in \{1, 2\}$. Let $G(X)$ be the resulted graph.

We have the following lemma on the chromatic number of $G(X)$.

Lemma 9 *The graph $G(X)$ is 4-colorable for each color class.*

We postpone the proof of this lemma until the end of this section and prove Lemma 1 first.

Proof of Lemma 1: By Lemma 8, there is a proper 133-coloring of G^* . We assume $V(G^*) = V(G) = \cup_{i=1}^{133} X_i$, where X_i is the i -th color class.

For each $i \in \{1, \dots, 133\}$, Lemma 9 shows $G(X_i)$ is 4-colorable; let $c_i: V(G(X_i)) \rightarrow T_i$ be a proper 4-coloring of the graph $G(X_i)$. Here $T_1, T_2, \dots, T_{133}$ are pairwise disjoint; each of them consists of 4 colors. For $i \in \{1, \dots, 133\}$, the 4-coloring c_i can be viewed as a 4-coloring of $G \setminus X_i$ since each vertex with label j receives the color $c_i(w_j)$ for $j = 1, 2$ and each removed vertex has at most three neighbors in $G \setminus X_i$.

Now we reuse the notation c_i to denote this 4-coloring of $G \setminus X_i$. For each $v \in X_i$, we have $|\cup_{u \in \Gamma_G(v)} c_i(u)| \leq 2$. We can assign two unused colors, denoted by the set $Y(v)$, to v . We define $f_i: V(G) \rightarrow \mathcal{P}(T_i)$ (the power set of T_i) satisfying

$$f_i(v) = \begin{cases} \{c_i(v)\} & \text{if } v \in V(G) \setminus X_i, \\ Y(v) & \text{if } v \in X_i. \end{cases}$$

Observe that each vertex in X_i receives two colors from f_i and every other vertex receives one color. Let $\sigma: V(G) \rightarrow \mathcal{P}(\cup_{i=1}^m T_i)$ be a mapping such that $\sigma(v) = \cup_{i=1}^m f_i(v)$. It is easy to verify σ is a 134-fold coloring of G such that each color is drawn from a palette of 532 colors; namely we have

$$\chi_f(G) \leq \frac{532}{134} = 4 - \frac{2}{67}.$$

The proof of Lemma 1 is finished. $\square$

Before we prove Lemma 9, we need the following definitions.

A *block* of a graph is a maximal 2-connected induced subgraph. A *Gallai tree* is a connected graph in which all blocks are either complete graphs or odd cycles. A *Gallai forest* is a graph all of whose components are Gallai trees. A *k-Gallai tree (forest)* is a Gallai

tree (forest) such that the degree of all vertices are at most $k - 1$. A k -critical graph is a graph G whose chromatic number is k and deleting any vertex can decrease the chromatic number. Gallai showed the following Lemma.

Lemma 10 [3] *If G is a k -critical graph, then the subgraph of G induced on the vertices of degree $k - 1$ is a k -Gallai forest.*

Proof of Lemma 9: We use proof by contradiction. Suppose that $G(X)$ is not 4-colorable. The only possible vertices in $G(X)$ with degree greater than 4 are the vertices w_1 and w_2 , which are obtained by contracting the vertices with label 1 and 2 in the intermediate graph H . The simple greedy algorithm shows that $G(X)$ is always 5-colorable. Let $G'(X)$ be a 5-critical subgraph of $G(X)$. Applying Lemma 10 to $G'(X)$, the subgraph of $G'(X)$ induced on the vertices of degree 4 is a 5-Gallai forest F . The vertex set of F may contain w_1 or w_2 . Delete w_1 and w_2 from F if F contains one of them. Let F' be the resulting Gallai forest. (Any induced subgraph of a Gallai forest is still a Gallai forest.) The Gallai forest F' is not empty. Let T be a connected component of F' and B be a leaf block of T . The block B is either a clique or an odd cycle from the definition of a Gallai tree.

Let v be a vertex in B . As v has at most two neighbors (w_1 and w_2) outside F' in $G(X)$, we have $d_{F'}(v) \geq 2$. If v is not in other blocks of F' , then we have $d_B(v) \geq 2$. It follows that $|B| \geq 3$. Since B is a subgraph of G and G is K_4 -free, the block B is an odd cycle.

Let $v_1 v_2$ be an edge in B such that v_1 and v_2 are not in other blocks. The degree requirement implies $v_i w_j$ are edges in $G(X)$ for all $i, j \in \{1, 2\}$. For $i = 1, 2$, there are vertices $x_i, y_i \in X$ satisfying $S(x_i) \cap \Gamma_G(v_i) \neq \emptyset$ and $S(y_i) \cap \Gamma_G(v_i) \neq \emptyset$; moreover either $|S(x_i)| = 4$ or $|S(y_i)| = 4$ since one of its neighborhood has label 2. Without loss of generality, we assume $|S(x_i)| = 4$ for $i \in \{1, 2\}$. If $x_i \neq y_i$, then $y_i \in N_G^4(x_i)$, i.e., $y_i \in \Gamma_{G^*}(x_i)$; this contradicts X being a color class. Thus we have $x_i = y_i$ and $|S(x_i)| = 4$ for $i \in \{1, 2\}$. For $\{i, j\} = \{1, 2\}$, if $x_i \neq y_j$, then $y_i \in N_G^5(x_i)$, i.e., $y_i \in \Gamma_{G^*}(x_i)$; this is a contradiction of X being a color class. Thus we have

$$x_1 = x_2 = y_1 = y_2.$$

Let x denote this common vertex above. Then $d_G(x) = 4$ and $\alpha(\Gamma_G(x)) = 2$.

Let v_0 be the only vertex in B shared by other blocks. Since $B - v_0$ is connected, the argument above shows there is a common x for all edges in $B - v_0$. If $\Gamma_{G(X)}(v_0) \cap \{w_1, w_2\} \neq \emptyset$, then there is some vertex $x_0 \in X$ such that $S(x_0) \cup \Gamma_G(v_0) \neq \emptyset$. By the similar argument, we also have $x_0 = x$.

Therefore, x depends only on B . In the sense that for any $y \in X$ and any $v \in B$, if $S(y) \cap \Gamma_G(v) \neq \emptyset$, then $y = x$.

The block B is an odd cycle as we mentioned above. Suppose $|B| = 2r + 1$. Let $v_0, v_1, \dots, v_{2r}$ be the vertices of B in cyclic order and v_0 be the only vertex which may be shared by other block.

Let $x \in X$ be the vertex determined by B . Recall $d_G(x) = 4$ and $\alpha(\Gamma_G(x)) = 2$. Each vertex in $\Gamma(x)$ can have at most 2 edges to B . We get

$$4r \leq |E(B, \Gamma(x))| \leq 8. \quad (4)$$

We have $r \leq 2$. The block B is either a C_5 or a K_3 . We claim both $v_0 w_1$ and $v_0 w_2$ are non-edges of $G(X)$.

If $B = C_5$, then inequality (4) implies that v_0 has no neighbor in $\Gamma(x)$ and the claim holds. If $B = K_3$, then the claim also holds; otherwise $B \cup \{S_1(x), S_2(x)\}$ forms a K_5^- in $G/S_1(x)/S_2(x)$, which is a contradiction to Lemma 6.

Let u_1 and u_2 be the two neighbors of v_0 in other blocks of F' . If u_1 and u_2 are in the same block, then this block is an odd cycle; otherwise, v_0u_1 and v_0u_2 are in two different blocks.

The union of non-leaf blocks of T is a Gallai-tree, denoted by T' . The argument above shows every leaf block of T' must be an odd cycle. Let C be such a leaf block of T' . Now C is an odd cycle, and C is connected to $|C| - 1$ leaf blocks of T . Let B and B' be two leaf blocks of T such that $B \cap C$ is adjacent to $B' \cap C$. Without loss of generality, we may assume B is the one we considered before. By the same argument, B' is an odd cycle of size $2r' + 1$ with $r' \in \{1, 2\}$. Let $v'_0, v'_1, \dots, v'_{2r'}$ be the vertices of B' and v'_0 be the only vertex in $B' \cap C$. For i in $\{1, 2, \dots, 2r'\}$ and j in $\{1, 2\}$, $v'_i w_j$ are edges in $G'(X)$. Similarly, there exists a vertex $x' \in X$ with $d_G(x') = 4$ and $\alpha(\Gamma_G(x')) = 2$ such that $|E(v_i, S_1(x'))| \geq 1$ and $|E(v_i, S_2(x'))| \geq 1$. We must have $x = x'$; otherwise $x' \in N_G^7(x)$, i.e., $x' \in \Gamma_{G^*}(x)$, and this contradicts the fact that X is a color class in D . Now we have $|E(\Gamma(x), B)| \geq 4r$ and $|E(\Gamma(x), B')| \geq 4r'$. By counting the degrees of vertices in $\Gamma(x)$ in G , we have

$$4r + 4r' + 4 + 4 \leq 16.$$

We get $r = r' = 1$. Both B and B' are K_3 's. In this case, $G/S_1(x)/S_2(x)$ contains the graph G_0 , see figure 9. This contradicts Lemma 6.

We can find the desired contradiction, so the lemma follows. $\square$

4 Proof of Lemma 6

In this section, we will prove Lemma 6. We first review a Lemma from [7].

Lemma 11 *Let G be a graph. Suppose that G_1 and G_2 are two subgraphs such that $G_1 \cup G_2 = G$ and $V(G_1) \cap V(G_2) = \{u, v\}$.*

1. *If uv is an edge of G , then we have*

$$\chi_f(G) = \max\{\chi_f(G_1), \chi_f(G_2)\}.$$

2. *If uv is not an edge of G , then we have*

$$\chi_f(G) \leq \max\{\chi_f(G_1), \chi_f(G_2 + uv), \chi_f(G_2/uv)\},$$

where $G_2 + uv$ is the graph obtained from G_2 by adding edge uv and G_2/uv is the graph obtained from G_2 by contracting $\{u, v\}$.

Proof of Lemma 6: Recall that G is a connected K_4 -free graph with minimum number of vertices such that $G \neq C_8^2$ and $\chi_f(G) > 4 - \frac{2}{67}$. Note that G is 2-connected. We will prove it by contradiction.

Suppose Lemma 6 fails for some vertex x in G . Observe $\Gamma_G(x)$ is one of the graphs in Figure 14. Here we assume $\Gamma_G(x) = \{a, b, c, d\}$. Through the proof of the lemma, let S_1 and S_2 be two vertex-disjoint independent sets in $\Gamma_G(x)$, H be a triangle in $V(G) \setminus (\{x\} \cup \Gamma_G(x))$, then say (S_1, S_2, H) is a bad triple if $\{\underline{S_1}, \underline{S_2}, H\}$ contains a K_5^- in $G/S_1/S_2$.

If $\Gamma_G(x) = P_4$, then $\{a, d\}$ and $\{b, c\}$ is the only pair of disjoint non-edges. There is a triangle H with $V(H) = \{y, z, w\}$ such that $(\{a, d\}, \{b, c\}, H)$ is a bad triple. Note that $|E(\{a, b, c, d\}, \{y, z, w\})| = 5$ or 6 . By an exhaustive search, the induced subgraph of G on $\{x, a, b, c, d, y, z, w\}$ is one of the following six graphs (see Figure 15).

If $\Gamma_G(x) = 2e$, then $(\{a, c\}, \{b, d\})$ and $(\{a, d\}, \{b, c\})$ are two pairs of disjoint non-edges. By considering the degrees of vertices in $\Gamma_G(x)$, there is only one triangle H with




Figure 14: Three possible cases of $\Gamma_G(x)$.




Figure 15: If $\Gamma_G(x) = P_4$, then there are six possible induced subgraphs.




Figure 16: If $\Gamma_G(x) = 2e$, then there are three possible induced subgraphs.

$V(H) = \{y, z, w\}$ such that $(\{a, c\}, \{b, d\}, H)$ and $(\{a, d\}, \{b, c\}, H)$ are two bad triples. By an exhaustive search, the induced subgraph of G on $\{x, a, b, c, d, y, z, w\}$ is one of the following three graphs (see Figure 16).

It suffices to show that G cannot contain H_i for $1 \leq i \leq 9$. Since all vertices in H_1 (and H_2) have degree 4, H_1 (and H_2) is the entire graph G . Observe that H_1 is isomorphic to C_8^2 and H_2 is 11:3-colorable (see Figure 17). Contradiction!




Figure 17: The graph H_2 and H_7 are 11:3-colorable.

In H_7 , the vertex d is the only vertex with degree less than 4. If H_7 is not the entire graph G , then d is a cut vertex of G . This contradicts the fact that G is 2-connected. Thus $G = H_7$. The graph H_7 is 11:3-colorable as shown by Figure 17. Contradiction!

Now we consider the case H_3 . Note $H_3 + bz$ is the graph H_2 . We have $\chi_f(H_3) \leq \chi_f(H_2) \leq 11/3$. The graph H_3 must be a proper induced subgraph of G , and the pair $\{b, z\}$ is a vertex cut of G . Let G' be the induced subgraph of G by deleting all vertices in H_3 but b, z . We apply Lemma 11 to $G + bz$ with $G_1 = H_3 + bz = H_2$ and $G_2 = G' + bz$. We have

$$\chi_f(G + bz) \leq \max\{\chi_f(H_2), \chi_f(G' + bz)\}.$$

Note $\chi_f(H_2) \leq 11/3$ and $11/3 < \chi_f(G) \leq \chi_f(G + bz)$. We have $\chi_f(G) \leq \chi_f(G' + bz)$. Both b and z have at most 2 neighbors in $G' + bz$. Thus $G' + bz$ is K_4 -free; $G' + bz \neq C_8^2$ and has fewer vertices than G . This contradicts to the minimality of G .

Note $H_5 + cy = H_2$. The case H_5 is similar to the case H_3 .

Note that H_4 , H_6 , and H_8 are isomorphic to each other. It suffices to show G does not contain H_4 . Suppose that H_4 is a proper induced subgraph of G . Let G_1 be the induced subgraph of G by deleting all vertices in H_4 . Note C_8^2 is not a proper subgraph of any graph in $\mathcal{G}_4$. We have $G_1 \neq C_8^2$. Note that c and z have degree 3 while other vertices in H_4 have degree 4. Since G is 2-connected, c has a unique neighbor, denoted by u , in $V(G_1)$. Similarly, z has a unique neighbor, denoted by v , in $V(G_1)$. Observe that the pair $\{u, v\}$ forms a vertex cut of G . Let G_2 be the induced graph of G on $V(H_4) \cup \{u, v\}$. Applying Lemma 11 to G with G_1 and G_2 , we have

$$\chi_f(G) \leq \max\{\chi_f(G_1), \chi_f(G_2 + uv), \chi_f(G_2/uv)\}.$$

Figure 18 shows $\chi_f(G_2 + uv)$ and $\chi_f(G_2/uv)$ are at most $11/3$.

Since $\chi_f(G) > 11/3$, we have $\chi_f(G) \leq \chi_f(G_1)$. Now G_1 is K_4 -free and has maximum degree at most 4; G_1 has fewer vertices than G . This contradicts the minimality of G .

If $G = H_4$, then $\chi_f(H_4) \leq 11/3$, since H_4 is a subgraph of $G_2 + uv$ in Figure 18.

Now we consider the last case H_9 . First, we contract b, c, z into a fat vertex denoted by $\underline{bcz}$. We write G/bcz for the graph after this contraction. Observe that $\{\underline{bcz}, d\}$ is a vertex-cut of G/bcz . Let G_4 and G'_4 be two connected subgraphs of G/bcz such that $G_4 \cup G'_4 = G/bcz$, $G_4 \cap G'_4 = \{\underline{bcz}, d\}$, and $\{u, v\} \subset G'_4$. Note that G_4 is 11:3 colorable, see Figure 19. Now by Lemma 11, we have

$$\chi_f(G/bcz) \leq \max\{\chi_f(G_4), \chi_f(G'_4)\}.$$




Figure 18: Case H_4 : both graph $G_2 + uv$ and G_2/uv are 11:3-colorable.

As $\{b, c, z\}$ is an independent set, each $a:b$ -coloring of G/bcz gives an $a:b$ -coloring of G , that is $\chi_f(G/bcz) \geq \chi_f(G) > 11/3$. The graph G_4 is 11:3-colorable; see Figure 19. Thus we have $\chi_f(G'_4) \geq \chi_f(G/bcz) \geq \chi_f(G)$. It is easy to check that G'_4 has maximum degree 4, K_4 -free, and it is not C_8^2 . Hence G'_4 must contain a K_4 . Otherwise, it contradicts the minimality of G .

Second, we contract $\{b, d, z\}$ into a fat vertex $\underline{bdz}$ and denote the graph by G/bcz . Let G_5 and G'_5 be two connected subgraphs of G/bdz such that $G_5 \cup G'_5 = G/bdz$, $G_5 \cap G'_5 = \{\underline{bdz}, c\}$, and $\{u, v\} \subset G'_5$. Note that G_5 is 11:3-colorable; see Figure 19. By a similar argument, G'_5 must contain a K_4 .




Figure 19: Case H_9 : the graphs G_4 , G_5 , and G_6 are 11:3-colorable.

The remaining case is that both G'_4 and G'_5 have a K_4 when we contract b and z . Since the original graph G is K_4 -free, the K_4 in G'_4 (and in G'_5) must contain the fat vertex $\underline{bcz}$ (or $\underline{bdz}$), respectively. Note that each of the four vertices b, c, d, z has at most one edge leaving H_9 . There must be a triangle uvp in G and these four outward edges are connected to some element of $\{u, v, p\}$. The graph G/bz must contain the subgraph G_6 as drawn in Figure 19.

Note that $\{u, v\}$ is a vertex-cut in G/bz . Let G_6 and G'_6 be two connected subgraphs of G/bz , which satisfy $G_6 \cup G'_6 = G$, $G_6 \cap G'_6 = \{u, v\}$, and $\underline{bz} \in G_6$. By Lemma 11, we have

$$\chi_f(G/bz) \leq \max\{\chi_f(G_6), \chi_f(G'_6)\}.$$

Note that G_6 is 11:3-colorable; see Figure 19. We also have $\chi_f(G/bz) \geq \chi_f(G) > \frac{11}{3}$. We obtain $\chi_f(G'_6) \geq \chi_f(G/bz) \geq \chi_f(G)$. Observe that G'_6 is a subgraph of G . We arrive at a contradiction of the minimality of G .

If $\Gamma_G(x) = C_4$, then the only possible choice for the two independent sets are $\{a, c\}$ and $\{b, d\}$. If there is some triangle H such that $(\{a, c\}, \{b, d\}, H)$ is a bad triple, then we have

$$|E(\Gamma_G(x), H)| \geq 5.$$

However, $|E(\Gamma_G(x), H)| \leq 4$. This is a contradiction. Thus the lemma follows in this case.

We can select two vertex disjoint non-edges S_1 and S_2 such that the graph $G/S_1/S_2$ contains no K_5^- . For these particular S_1 and S_2 , if $G/S_1/S_2$ contains no G_0 , then Lemma 6 holds.

Without loss of generality, we assume that $G/S_1/S_2$ does contain G_0 . Let $s_i = \underline{S_i}$ for $i = 1, 2$. Observe that both s_1 and s_2 have four neighbors u, v, p, q other than x in $\underline{G_0}$. It follows that

$$|E(S_1 \cup S_2, \{u, v, p, q\})| \geq 8.$$

On the one hand, we have

$$\begin{aligned} |E(G \mid_{S_1 \cup S_2})| &= \frac{1}{2} \left(\sum_{v \in S_1 \cup S_2} d(v) - |E(S_1 \cup S_2, \{u, v, p, q\})| - 4 \right) \\ &\leq \frac{1}{2} (16 - 8 - 4) \\ &= 2. \end{aligned}$$

On the other hand, $\alpha(\Gamma(x)) = 2$ implies $G \mid_{S_1 \cup S_2}$ contains at least two edges. Thus, we have $\Gamma_G(x) = 2$ e. Label the vertices in $\Gamma_G(x)$ by a, b, c, d as in Figure 14. We assume ab and cd are edges while ac, bd, ad, bc are non-edges. Observe that each vertex in $\{u, v, p, q\}$ has exactly two neighbors in $\{a, b, c, d\}$.

If one vertex, say u , has two neighbors forming a non-edge, say ac , then we can choose $S'_1 = \{a, c\}$ and $S'_2 = \{b, d\}$. It is easy to check that $G/S'_1/S'_2$ contains neither G_0 nor K_5^- . We are done in this case.

In the remaining case, we can assume that for each vertex y in $\{u, v, p, q\}$, the neighbors of y in $\{a, b, c, d\}$ always form an edge. Up to relabeling vertices, there is only one arrangement for edges between $\{u, v, p, q\}$ and $\{a, b, c, d\}$; see the graph H_{10} defined in Figure 20. The




Figure 20: H_{10} and an 11:3-coloring of H_{10} .

graph H_{10} is 11:3-colorable as shown in Figure 20. Since $\chi_f(G) > 11/3$, H_{10} is a proper subgraph of G . Note in H_{10} , every vertices except w and r has degree 4; both w and r have degree 3. Thus, $\{w, r\}$ is a vertex cut of G . Let $G_1 = H_{10}$ and G_2 be the subgraph of G by deleting vertices in $\{x, a, b, c, d, p, q, u, v\}$. Applying Lemma 11 with G_1 and G_2 defined above, we have

$$\chi_f(G) \leq \max\{\chi_f(G_1), \chi_f(G_2)\}.$$

Since $\chi_f(G) > 11/3$ and $\chi_f(G_1) \leq 11/3$ (see Figure 20), we must have $\chi_f(G_2) \geq \chi_f(G)$. Note that G_2 has fewer number of vertices than G . This contradicts the minimality of G . Therefore, the lemma follows. $\square$

References

- [1] O. Borodin and A. Kostochka, On an upper bound on a graph's chromatic number, depending on the graph's degree and density, *J. Comb. Theory Ser. B*, **23** (1977), 247-250.
- [2] D. Christofides, K. Edwards, and A.D. King, A note on hitting maximum and maximal cliques with a stable set, *Submitted. Arxiv preprint 1109.3092*, (2011).
- [3] T. Gallai, Kritische graphen I, *Magyar Tud. Akad. Mat. Kutat ó Int. Közl*, **8** (1963), 165-192.
- [4] H. Hatami and X. Zhu, The fractional chromatic number of graphs of maximum degree at most three, *SIAM. J. Discrete Math.*, **24** (2009), 1762-1775.
- [5] C. C. Heckman and R. Thomas, A new proof of the independence ratio of triangle-free cubic graphs, *Discrete Math.*, **233** (2001), 233-237.
- [6] Andrew D. King, Hitting all maximum cliques with a stable set using lopsided independent transversals, *J. Graph Theory*, **67-4** (2011), 300-305.
- [7] L. Lu and X. Peng, The fractional chromatic number of triangle-free graphs with $\Delta \leq 3$, *submitted*.
- [8] M. Molloy and B. Reed, *Graph colouring and the probabilistic method*, volume 23 of Algorithms and Combinatorics, Springer-Verlag, Berlin, 2002.
- [9] B. Reed, ω , Δ , and χ , *J. Graph Theory*, **27-4** (1998), 177-227.
- [10] B. Reed, A strengthening of Brooks' theorem, *J. Comb. Theory Ser. B*, **76** (1999), 136-149.
- [11] E. R. Scheinerman and D. H. Ullman, *Fractional graph theory*, A rational approach to the theory of graphs, Wiley-Intersci. Ser. Discrete Math. Optim, John Wiley & Sons, Inc, New York, 1997.

A Fractional Analogue of Brooks' Theorem

Andrew D. King ^{*} Linyuan Lu [†] Xing Peng [‡]

January 15, 2013

Abstract

Let $\Delta(G)$ be the maximum degree of a graph G . Brooks' theorem states that the only connected graphs with chromatic number $\chi(G) = \Delta(G) + 1$ are complete graphs and odd cycles. We prove a fractional analogue of Brooks' theorem in this paper. Namely, we classify all connected graphs G such that the fractional chromatic number $\chi_f(G)$ is at least $\Delta(G)$. These graphs are complete graphs, odd cycles, C_8^2 , $C_5 \boxtimes K_2$, and graphs whose clique number $\omega(G)$ equals the maximum degree $\Delta(G)$. Among the two sporadic graphs, the graph C_8^2 is the square graph of cycle C_8 while the other graph $C_5 \boxtimes K_2$ is the strong product of C_5 and K_2 . In fact, we prove a stronger result; if a connected graph G with $\Delta(G) \geq 4$ is not one of the graphs listed above, then we have $\chi_f(G) \leq \Delta(G) - \frac{2}{67}$.

1 Introduction

The chromatic number of graphs with bounded degrees has been studied for many years. Brooks' theorem perhaps is one of the most fundamental results; it is included by many textbooks on graph theory. Given a simple connected graph G , let $\Delta(G)$ be the maximum degree, $\omega(G)$ be the clique number, and $\chi(G)$ be the chromatic number. Brooks' theorem states that $\chi(G) \leq \Delta(G)$ unless G is a complete graph or an odd cycle. Reed [9] proved that $\chi(G) \leq \Delta(G) - 1$ if $\omega(G) \leq \Delta(G) - 1$ and $\Delta(G) \geq \Delta_0$ for some large constant Δ_0 . This excellent result was proved by probabilistic methods, and Δ_0 is at least hundreds. Before this result, Borodin and Kostochka [1] made the following conjecture.

Conjecture [1]: Suppose that G is a connected graph. If $\omega(G) \leq \Delta(G) - 1$ and $\Delta(G) \geq 9$, then we have

$$\chi(G) \leq \Delta(G) - 1.$$

If the conjecture is true, then it is best possible since there is a K_8 -free graph $G = C_5 \boxtimes K_3$ (actually K_7 -free, see Figure 1) with $\Delta(G) = 8$ and $\chi(G) = 8$.

Here we use the following notation of the strong product. Given two graphs G and H , the *strong product* $G \boxtimes H$ is the graph with vertex set $V(G) \times V(H)$, and (a, x) is connected to (b, y) if one of the following holds

- $a = b$ and $xy \in E(H)$,
- $ab \in E(G)$ and $x = y$,

^{*}Simon Fraser University, Burnaby, British Columbia, Canada (andrew.d.king@gmail.com).

[†]University of South Carolina, Columbia, SC 29208, (lu@math.sc.edu). This author was supported in part by NSF grant DMS 1000475.

[‡]University of South Carolina, Columbia, SC 29208, (pengx@mailbox.sc.edu). This author was supported in part by NSF grant DMS 1000475.




Figure 1: The graph $C_5 \square K_3$.

- $ab \in E(G)$ and $xy \in E(H)$.

Reed's result [9] settled Borodin and Kostochka's conjecture for sufficiently large $\Delta(G)$, but the cases with small $\Delta(G)$ are hard to cover using the probabilistic method.

In this paper we consider a fractional analogue of this problem. The fractional chromatic number $\chi_f(G)$ can be defined as follows. A b -fold coloring of G assigns a set of b colors to each vertex such that any two adjacent vertices receive disjoint sets of colors. We say a graph G is $a:b$ -colorable if there is a b -fold coloring of G in which each color is drawn from a palette of a colors. We refer to such a coloring as an $a:b$ -coloring. The b -fold coloring number, denoted by $\chi_b(G)$, is the smallest integer a such that G has an $a:b$ -coloring. Note that $\chi_1(G) = \chi(G)$. It was shown that $\chi_{a+b}(G) \leq \chi_a(G) + \chi_b(G)$. The fractional chromatic number $\chi_f(G)$ is $\lim_{b \rightarrow \infty} \frac{\chi_b(G)}{b}$.

By the definition, we have $\chi_f(G) \leq \chi(G)$. The fractional chromatic number can be viewed as a relaxation of the chromatic number. Many problems involving the chromatic number can be asked again using the fractional chromatic number. The fractional analogue often has a simpler solution than the original problem. For example, the famous $\omega - \Delta - \chi$ conjecture of Reed [8] states that for any simple graph G , we have

$$\chi(G) \leq \left\lceil \frac{\omega(G) + \Delta(G) + 1}{2} \right\rceil.$$

The fractional analogue of $\omega - \Delta - \chi$ conjecture was proved by Molloy and Reed [7]; they actually proved a stronger result with ceiling removed, i.e.,

$$\chi_f(G) \leq \frac{\omega(G) + \Delta(G) + 1}{2}. \quad (1)$$

In this paper, we classify all connected graphs G with $\chi_f(G) \geq \Delta(G)$.

Theorem 1 *A connected graph G satisfies $\chi_f(G) \geq \Delta(G)$ if and only if G is one of the following*

1. a complete graph,
2. an odd cycle,
3. a graph with $\omega(G) = \Delta(G)$,

4. C_8^2 ,

5. $C_5 \boxtimes K_2$.

For the complete graph K_n , we have $\chi_f(K_n) = n$ and $\Delta(K_n) = n - 1$. For the odd cycle C_{2k+1} , we have $\chi_f(C_{2k+1}) = 2 + \frac{1}{k}$ and $\Delta(C_{2k+1}) = 2$. If G is neither a complete graph nor an odd cycle but contains a clique of size $\Delta(G)$, then we have

$$\Delta(G) \leq \omega(G) \leq \chi_f(G) \leq \chi(G) \leq \Delta(G). \quad (2)$$

The last inequality is from Brooks' theorem. The sequence of inequalities above implies $\chi_f(G) = \Delta(G)$.

If G is a vertex-transitive graph, then we have [10]

$$\chi_f(G) = \frac{|V(G)|}{\alpha(G)},$$

where $\alpha(G)$ is the independence number of G . Note that both graphs C_8^2 and $C_5 \boxtimes K_2$ are vertex-transitive and have the independence number 2. Thus we have

$$\chi_f(C_8^2) = 4 = \Delta(C_8^2) \quad \text{and} \quad \chi_f(C_5 \boxtimes K_2) = 5 = \Delta(C_5 \boxtimes K_2).$$



Figure 2: The graph C_8^2 and $C_5 \boxtimes K_2$.

Actually, Theorem 1 is a corollary of the following stronger result.

Theorem 2 *Assume that a connected graph G is neither C_8^2 nor $C_5 \boxtimes K_2$. If $\Delta(G) \geq 4$ and $\omega(G) \leq \Delta(G) - 1$, then we have*

$$\chi_f(G) \leq \Delta(G) - \frac{2}{67}.$$

Remark: In the case $\Delta(G) = 3$, Heckman and Thomas [4] conjectured that $\chi_f(G) \leq 14/5$ if G is triangle-free. Hatami and Zhu [3] proved $\chi_f(G) \leq 3 - \frac{3}{64}$ for any triangle-free graph G with $\Delta(G) \leq 3$. The authors showed an improved result $\chi_f(G) \leq 3 - \frac{3}{43}$ in the previous paper [6]. Thus we need only consider the cases $\Delta(G) \geq 4$. For any connected graph G with sufficiently large $\Delta(G)$ and $\omega(G) \leq \Delta(G) - 1$, Reed's result [9] $\chi(G) \leq \Delta(G) - 1$ implies $\chi_f(G) \leq \Delta(G) - 1$. The method introduced in [3] and strengthened in [6], has a strong influence on this paper. The readers are encouraged to read these two papers [3, 6].

Let $f(k) = \inf_G \{\Delta(G) - \chi_f(G)\}$, where the infimum is taken over all connected graphs G with $\Delta(G) = k$ and not one of the graphs listed in Theorem 1. Since $\chi_f(G) \geq \omega(G)$, by taking a graph with $\omega(G) = \Delta(G) - 1$, we have $f(k) \leq 1$. Theorem 2 says $f(k) \geq \frac{2}{67}$ for any $k \geq 4$. Reed's result [9] implies $f(k) = 1$ for sufficiently large k . Heckman and Thomas [4]

conjectured $f(3) = 1/5$. It is an interesting problem to determine the value of $f(k)$ for small k . Here we conjecture $f(4) = f(5) = \frac{1}{3}$. If Borodin and Kostochka's conjecture is true, then $f(k) = 1$ for $k \geq 9$.

Theorem 2 is proved by induction on k . Because the proof is quite long, we split the proof into the following two lemmas.

Lemma 1 *We have $f(4) \geq \frac{2}{67}$.*

Lemma 2 *For each $k \geq 6$, we have $f(k) \geq \min\{f(k-1), \frac{1}{2}\}$. We also have $f(5) \geq \min\{f(4), \frac{1}{3}\}$.*

It is easy to see the combination of Lemma 1 and Lemma 2 implies Theorem 2. The idea of reduction comes from the first author, who pointed out $f(k) \geq \min\{f(k-1), \frac{1}{2}\}$ for $k \geq 7$ based on his recent results [5]. The second and third authors originally proved $f(k) \geq \frac{C}{k^5}$ (for some $C > 0$) using different method in the first version; they also prove the reductions at $k = 5, 6$, which are much harder than the case $k \geq 7$. We do not know whether a similar reduction exists for $k = 4$.

The rest of this paper is organized as follows. In section 2, we will introduce some notation and prove Lemma 2. In section 3 and section 4, we will prove $f(4) \geq \frac{2}{67}$.

2 Proof of Lemma 2

In this paper, we use the following notation. Let G be a simple graph with vertex set $V(G)$ and edge set $E(G)$. The *neighborhood* of a vertex v in G , denoted by $\Gamma_G(v)$, is the set $\{u : uv \in E(G)\}$. The *degree* $d_G(v)$ of v is the value of $|\Gamma_G(v)|$. The *independent set* (or *stable set*) is a set S such that no edge with both ends in S . The *independence number* $\alpha(G)$ is the largest size of S among all the independent sets S in G . When $T \subset V(G)$, we use $\alpha_G(T)$ to denote the independence number of the induced subgraph of G on T . Let $\Delta(G)$ be the maximum degree of G . For any two vertex-sets S and T , we define $E_G(S, T)$ as $\{uv \in E(G) : u \in S \text{ and } v \in T\}$. Whenever G is clear under context, we will drop the subscript G for simplicity.

If S is a subset of vertices in G , then *contracting* S means replacing vertices in S by a single fat vertex, denoted by $\underline{S}$, whose incident edges are all edges that were incident to at least one vertex in S , except edges with both ends in S . The new graph obtained by contracting S is denoted by G/S . This operation is also known as *identifying vertices of* S in the literature. For completeness, we allow S to be a single vertex or even the empty set. If S only consists of a single vertex, then $G/S = G$; if $S = \emptyset$, then G/S is the union of G and an isolated vertex. When S consists of 2 or 3 vertices, for convenience, we write G/uv for $G/\{u, v\}$ and G/uvw for $G/\{u, v, w\}$; the fat vertex will be denoted by $\underline{uv}$ and $\underline{uvw}$, respectively. Given two disjoint subsets S_1 and S_2 , we can contract S_1 and S_2 sequentially. The order of contractions does not matter; let $G/S_1/S_2$ be the resulted graph. We use $G - S$ to denote the subgraph of G induced by $V(G) - S$.

In order to prove Lemma 2, we need use the following theorems due to King [5].

Theorem 3 (King [5]) *If a graph G satisfies $\omega(G) > \frac{2}{3}(\Delta(G) + 1)$, then G contains a stable set S meeting every maximum clique.*

Theorem 4 (King [5]) *For a positive integer k , let G be a graph with vertices partitioned into cliques $V_1, \dots, V_r$. If for every i and every $v \in V_i$, v has at most $\min\{k, |V_i| - k\}$ neighbors outside V_i , then G contains a stable set of size r .*

Lemma 3 Suppose that G is a connected graph with $\Delta(G) \leq 6$ and $\omega(G) \leq 5$. Then there exists an independent set meeting all induced copies of K_5 and $C_5 \boxtimes K_2$.

Proof: We first show that there exists an independent set meeting all copies of K_5 . If G contains no K_5 , then this is trivial. Otherwise, we can apply Theorem 3 to get the desired independent set since $\omega(G) > \frac{2}{3}(\Delta(G) + 1)$ is satisfied.

Now we prove the Lemma by contradiction. Suppose the Lemma is false. Let G be a minimum counterexample (with the smallest number of vertices). For any independent set I , let $C(I)$ be the number of induced copies of $C_5 \boxtimes K_2$ in $G - I$. Among all independent sets which meet all copies of K_5 , there exists one such independent set I such that $C(I)$ is minimized.

Since $C(I) > 0$, there is an induced copy of $C_5 \boxtimes K_2$ in $G - I$; we use H to denote it. In $C_5 \boxtimes K_2$, there is a unique perfect matching such that identifying the two ends of each edge in this matching results a C_5 . An edge in this unique matching is called a *canonical* edge. We define a new graph G' as follows: First we contract all canonical edges in H to get a C_5 , where its vertices are called *fat* vertices. Second we add five edges turning the C_5 into a K_5 . Observe that each vertex in this C_5 can have at most two neighbors in $G - H$ and $\Delta(G') \leq 6$. We will consider the following four cases.

Case 1: There is a K_6 in the new graph G' . Since the original graph G is K_6 -free, the K_6 is formed by the following two possible ways.

Subcase 1a: This K_6 contains 5 fat vertices. By the symmetry of H , there is an induced C_5 in H such that the vertices in C_5 contain a common neighbor vertex v in $G \setminus V(H)$, see Figure 3. Since H is K_5 -free, we can find x, y in this C_5 such that x, y is a non-edge. Let $I' := (I \setminus \{v\}) \cup \{x, y\}$; I' is also an independent set. Observe that v is not in any $K_5 \subset G - I'$. Thus the set I' is also an independent set and meets every K_5 in G . Since $C_5 \boxtimes K_2$ is a 5-regular graph, any copy of $C_5 \boxtimes K_2$ containing v must contain at least one of x and y . Thus, $C(I') < C(I)$. Contradiction!




Figure 3: Subcase 1a.




Figure 4: Subcase 1b.

Subcase 1b: This K_6 contains 4 fat vertices. Let u, v be the other two vertices. By the symmetry of H , there is a unique way to connect u and v to H as shown by Figure 4. Since uv is an edge, one of u and v is not in I . We assume $u \notin I$. Let $\{x, y\} \subset \Gamma_G(v) \cap V(H)$ as shown in Figure 4 and $I' = I \setminus \{v\} \cup \{x, y\}$. Observe that I' is an independent set and $v \notin K_5 \subset G - I'$. Thus I' is an independent set meeting each K_5 in G . Since each $C_5 \boxtimes K_2$ containing v must contain one of x and y . Thus $C(I') < C(I)$. Contradiction!

Case 2: There is a K_5 intersecting H with 4 vertices. Let v be the vertex of this K_5 but not in H , see Figure 5. We have two subcases.

Subcase 2a: The vertex v has another neighbor y in H but not in this K_5 . Since H is K_5 -free, we can select a vertex x in this K_5 such that xy is not an edge of G . Let $I' := I \setminus \{v\} \cup \{x, y\}$. Note that $v \notin K_5 \subset G - I'$ and I' is an independent set. Thus I' is an independent set meeting each K_5 in G . Since any $C_5 \boxtimes K_2$ containing v must contain one of x and y , we have $C(I') < C(I)$. Contradiction!

Subcase 2b: All neighbors of v in H are in this K_5 . Let x be any vertex in this K_5 other than v , and $I' := I \setminus \{v\} \cup \{x\}$. In this case, there is only one K_5 containing v . Thus, I' is also an independent set meeting every copy of K_5 in G . Observe that $\Gamma_G(v) \setminus \{x\}$ is disconnected. If $v \in H' = C_5 \boxtimes K_2$, then $\Gamma_G(v) \cap H'$ is connected. Thus $v \notin C_5 \boxtimes K_2 \subset G - I'$ and $C(I') < C(I)$. Contradiction!




Figure 5: Case 2.




Figure 6: Case 3.

Case 3: There is an induced subgraph H' isomorphic to $C_5 \boxtimes K_2$ such that H' and H are intersecting, see Figure 6. Since $V(H) \cap V(H') \neq \emptyset$ and $H \neq H'$, we can find a canonical edge uv of H and a canonical edge uv' of H' such that $v \notin V(H')$ and $v' \notin V(H)$. If vv' is a non-edge, then let $I' := I \setminus \{v'\} \cup \{u\}$. It is easy to check I' is still an independent set. We also observe that any possible K_5 containing v' must also contain u . Thus, I' meets every copy of K_5 in G . We have $v' \notin C_5 \boxtimes K_2 \subset G - I'$ since vv' is not an edge. We get $C(I') < C(I)$. Contradiction! If vv' is an edge, then locally there are two K_5 intersecting at u, v , and v' ; say the other four vertices are x_1, x_2, y_1, y_2 , where two cliques are $\{x_1, x_2, u, v, v'\}$ and $\{y_1, y_2, u, v, v'\}$, see Figure 6. Let $I' = I \cup \{x_1, y_1\} \setminus \{v'\}$. Note that I' is an independent set and $v' \notin K_5 \subset G - I'$. Thus I' is an independent set meeting each K_5 in G . Observe that any copies of $C_5 \boxtimes K_2$ containing v' must contain one of x_1 and y_1 ; we have $C(I') < C(I)$. Contradiction!

Case 4: This is the remaining case, G' is K_6 -free. We have $\omega(G') \leq 5$ and $|V(G')| < |V(G)|$. By the minimality of G , there is an independent set I' of G' meeting every copy of K_5 and $C_5 \boxtimes K_2$. In I' , there is a unique vertex x of the K_5 obtained from contracting canonical edges of H . Let uv be the canonical edge corresponding to x . Let $I'' = I' \setminus \{x\} \cup \{u\}$, we get an independent set I'' of G . Note that any $v \in H \setminus \{u\}$ is not in any K_5 of $G - I''$ by Case 2 as well as not in any $C_5 \boxtimes K_2$ of $G - I''$ by Case 3. Thus I'' hits each K_5 in G and $C(I'') = 0$. Contradiction! $\square$

Lemma 4 *Let G be a connected graph with $\Delta(G) \leq 5$ and $\omega(G) \leq 4$. If $G \neq C_{2l+1} \boxtimes K_2$ for some $l \geq 2$, then there is an independent set I hitting all copies of K_4 in G .*

Proof: We will prove it by contradiction. If the lemma is false, then let G be a minimum counterexample. If G is K_4 -free, then there is nothing to prove. Otherwise, we consider the clique graph $\mathcal{C}(G)$, whose edge set is the set of all edges appearing in some copy of K_4 . Because of $\Delta(G) = 5$, here are all possible connected component of $\mathcal{C}(G)$.

1. $C_t \boxtimes K_2$ for $t \geq 4$. If this type occurs, then every vertex in $C_t \boxtimes K_2$ has degree 5; thus, this is the entire graph G . If t is even, then we can find an independent set I meeting every K_4 . If t is odd, then it is impossible to find such an independent set. However, this graph is excluded from the assumption of the Lemma.
2. $P_t \boxtimes K_2$ for $t \geq 3$. In this case, all internal vertices have degree 5 while the four end vertices have degree 4. Consider a new graph G' which is obtained by deleting all

internal vertices and adding four edges to make the four end vertices as a K_4 . It is easy to check $\Delta(G') \leq 5$ and $\omega(G') \leq 4$. Since $|G'| < |G|$, there is an independent set I of G' meeting every copy of K_4 in G' . Note that there is exactly one end vertex in I . Observe that any one end vertex can be extended into a maximal independent set meeting every copy of K_4 in $P_t \boxtimes K_2$. Thus, we can extend I to an independent set I' of G such that I' meets every copy of K_4 in G . Hence, this type of component does not occur in $\mathcal{C}(G)$.

3. There are four other types listed in Figure 7.




Figure 7: All types of components in the clique graph $\mathcal{C}(G)$.

For each component C_i in $\mathcal{C}(G)$, let V_i be the set of common vertices in all K_4 's of C_i ; for the leftmost figure in Figure 7, V_i is the set of all 4 vertices; for the middle two figures, V_i is the set of bottom three vertices; for the rightmost figure, V_i consists of the left-bottom vertex and the middle-bottom vertex. Note that all V_i 's are pairwise disjoint. Let G' be the induce subgraph of G on $\cup_i V_i$. Note that G' does not contains any vertex in $C_i \setminus V_i$. By checking each type, we find out that for each i and each $v \in V_i$, v has at most $\min\{2, |V_i| - 2\}$ neighbors outside V_i in G' (not in G !). Applying Theorem 4 to G' , we conclude that there exists an independent set I of G' meeting every V_i ; thus I meets every K_4 in G . Contradiction!

□

Lemma 5 *Let G be a connected graph with $\Delta(G) \leq 5$ and $\omega(G) \leq 4$. If $G \neq C_{2l+1} \boxtimes K_2$ for some $l \geq 2$, then there exists an independent set meeting all induced copies of K_4 and C_8^2 .*

Proof: We will use proof by contradiction. Suppose the Lemma is false. Let G be a minimum counterexample (with the smallest number of vertices). For any independent set I , let $C(I)$ be the number of induced copies of C_8^2 in $G - I$. Among all independent sets which meet all copies of K_4 , there exists an independent set I such that $C(I)$ is minimized. Since $C(I) > 0$, let H be a copy of C_8^2 in $G - I$. The vertices of H are listed by u_i for $i \in \mathbb{Z}_8$ anticlockwise such that $u_i u_j$ is an edge of H if and only if $|i - j| \leq 2$. The vertex v_{i+4} is the *antipode* of v_i for any $i \in \mathbb{Z}_8$.

Case 1: There exists a vertex $v \notin V(H)$ such that v has five neighbors in H . By the Pigeonhole Principle, $\Gamma(v)$ contains a pair of antipodes. Without loss of generality, say $u_0, u_4 \in \Gamma(v)$. If the other three neighbors of v do not form a triangle, then we let $I' := I \setminus \{v\} \cup \{u_0, u_4\}$; note that v is not in any K_4 of $G - I'$. Thus I' is an independent set meeting every copy of K_4 . Since every copy of C_8^2 containing v must contain one of u_0 and u_4 , we have $C(I') < C(I)$. Contradiction! Hence, the other three neighbors of v must form a triangle. Without loss of generality, we can assume that the three neighbors are u_1, u_2 , and u_3 . Now we let $I'' := I \setminus \{v\} \cup \{u_0, u_3\}$; note that $v \notin K_4 \subset G - I'$. Thus I'' is also an independent set meeting every copy of K_4 of G . Since every copy of C_8^2 containing v must contain one of u_0 and u_3 , we have $C(I'') < C(I)$. Contradiction!

Case 2: There exists a vertex $v \notin V(H)$ such that v has exactly four neighbors in H . Since H is K_4 -free, we can find $u_i, u_j \in \Gamma(v) \cap V(H)$ such that $u_i u_j$ is a non-edge. Let

$I' := I \setminus \{v\} \cup \{u_i, u_j\}$; I' is also an independent set. Note that $\Gamma(v) \setminus \{u_i, u_j\}$ can not be a triangle, v is not in any $K_4 \subset G - I'$. Thus I' meets every copy of K_4 . Since every copy of C_8^2 containing v must contain one of u_i and u_j , we have $C(I') < C(I)$. Contradiction!

Case 3: There exists a vertex $v \notin V(H)$ such that v has exactly three neighbors in H . If the 3 neighbors do not form a triangle, then choose $u_i, u_j \in \Gamma(v) \cap V(H)$ such that $u_i u_j$ is a non-edge. Note that $\Gamma(v) \setminus \{u_i, u_j\}$ can not be a triangle; v is not in any $K_4 \subset G - I'$. Let $I' := I \setminus \{v\} \cup \{u_i, u_j\}$; I' is also an independent set meeting every copy of K_4 . Since every copy of C_8^2 containing v must contain one of u_i and u_j , we have $C(I') < C(I)$. Contradiction! Else, the three neighbors form a triangle; let u_i be one of them and $I' := I \setminus \{v\} \cup \{u_i\}$; v is not in any $K_4 \subset G - I'$. Thus I' is an independent set meeting every copy of K_4 . Note that $\Gamma(v) \setminus \{u_i\}$ has only two vertices in H . The induced graph on $\Gamma(v) \setminus \{u_i\}$ is disconnected. However, for any vertex v in $H' = C_8^2$, the subgraph induced by $\Gamma_G(v) \cap H'$ is a P_4 . There is no C_8^2 in $G - I'$ containing v . Thus, $C(I') < C(I)$. Contradiction!

Case 4: Every vertex outside H can have at most 2 neighbors in H . We identify each pair of antipodes of H to get a new graph G' from G . After identifying, H is turned into a K_4 ; where the vertices of this K_4 are referred as fat vertices.

Subcase 4a: $G' \neq C_{2l+1} \boxtimes K_2$. Observe $\Delta(G') \leq 5$. We claim G' is K_5 -free. Suppose not. Since every vertex in H has at most one neighbor outside H , then each fat vertex can have at most two neighbors outside H . Recall that the original graph G is K_5 -free. If G' has some K_5 , then this K_5 contains either 3 or 4 fat vertices. Let w be one of the other vertices in this K_5 . We get w has at least three neighbors in H . However, this is covered by Case 1, Case 2, or Case 3. Thus, G' is K_5 -free. Since $|G'| < |G|$, by the minimality of G , G' has an independent set I meeting every copy of K_4 and C_8^2 in G' . There is exactly one fat vertex in I . Now replacing this fat vertex by its corresponding pair of antipodal vertices, we get an independent set I' ; we assume the pair of antipodal vertices are u_2 and u_6 . It is easy to check that I' is an independent set of G . Next we claim any $v \in V(H) \setminus \{u_2, u_6\}$ is neither in a $K_4 \subset V(G) - I'$ nor in a $C_8^2 \subset V(G) - I'$. Suppose there is some v such that $v \in K_4 \subset G - I'$. Recall each $v \in V(H)$ has at most one neighbor outside H and H is K_4 -free; there is some $w \notin V(H)$ such that w has at least three neighbors in H . This is already considered by Case 1, Case 2, or Case 3. We are left to show that $v \notin C_8^2 \subset G - I'$ for each $v \in V(H) \setminus \{u_2, u_6\}$. If not, there exists a copy H' of C_8^2 in $G - I'$ containing v . Note H' is 4-regular, any vertex in H' can have at most one neighbor in I' ; in particular, $v \neq u_0, u_4$. Without loss of generality, we assume $v = u_3$. Then there is a vertex $w \notin V(H)$ such that $u_3 w$ is an edge, see Figure 8. Observe that the neighborhood of each vertex of an induced C_8^2 is is a P_4 . Since $u_1 u_4$ and $u_1 u_5$ are two non-edges, we have $w u_1$ being an edge. Observe $\Gamma_G(u_1) = \{u_7, u_0, u_2, u_3, w\}$. Since $u_2 \notin H'$, we have $u_0 \in H'$; u_0 has two neighbors (u_2 and u_6) outside H' , contradiction! Therefore, I' meets every copy of K_4 and C_8^2 in G . Contradiction!




Figure 8: Subcase 4a.

Subcase 4b: $G' = C_{2l+1} \boxtimes K_2$. The graph G can be recovered from G' . It consists of an induced subgraph $H = C_8^2$ and an induced subgraph $P_{2l-1} \boxtimes K_2$. For each vertex u in H , there is exactly one edge connecting it to one of the four end vertices of $P_{2l-1} \boxtimes K_2$; for each end vertex v of $P_{2l-1} \boxtimes K_2$, there are exactly two edges connecting v to the vertices in H . First, we take any maximum independent set I' of $P_{2l-1} \boxtimes K_2$. Observe that I' has exactly two end points of $P_{2l-1} \boxtimes K_2$; so I' has exactly four neighbors in H . In the remaining four vertices of H , there exists a non-edge $u_i u_j$ since H is K_4 -free. Let $I := I' \cup \{u_i, u_j\}$. Clearly I is an independent set of G meeting every copy of K_4 and C_8^2 . Contradiction! $\square$

We are ready to prove Lemma 2.

Proof of Lemma 2: We need prove for $k \geq 5$ and any connected graph G with $\Delta(G) = k$ and $\omega(G) \leq k - 1$ satisfies

$$\chi_f(G) \leq k - \min \left\{ f(k-1), \frac{1}{2} \right\}. \quad (3)$$

If $\omega(G) \leq k - 2$, then by inequality (1), we have

$$\chi_f(G) \leq \frac{\Delta(G) + \omega(G) + 1}{2} \leq k - \frac{1}{2}.$$

Thus, inequality (3) is satisfied. From now on, we assume $\omega(G) = \Delta(G) - 1$.

For $\Delta(G) = k \geq 6$ and $\omega(G) = k - 1$, the condition $\omega(G) > \frac{2}{3}(\Delta(G) + 1)$ is satisfied. By Theorem 3, G contains an independent set meeting every maximum clique. Extend this independent set to a maximal independent set and denote it by I . Note that $\Delta(G - I) \leq k - 1$ and $\omega(G - I) \leq k - 2$.

Case 1: $k \geq 7$. From the definition of $f(k - 1)$, we have $\chi_f(G - I) \leq \Delta(G - I) - f(k - 1)$. Thus,

$$\chi_f(G) \leq \chi_f(G - I) + 1 \leq k - 1 - f(k - 1) + 1 = k - f(k - 1).$$

Thus, we have $f(k) \geq \min\{f(k - 1), 1/2\}$.

Case 2: $k = 6$. By Lemma 3, we can find an independent set meets every copy of K_5 and $C_5 \boxtimes K_2$; we extend this independent set as a maximal independent set I . Note that $G - I$ contains no induced subgraph isomorphic $C_5 \boxtimes K_2$. We have $\chi_f(G - I) \leq 5 - f(5)$; it implies $\chi_f(G) \leq 6 - f(5)$. Thus, $f(6) \geq \min\{f(5), 1/2\}$ and we are done.

Case 3: $k = 5$. If $G = C_{2l+1} \boxtimes K_2$ for some $l \geq 3$; then G is vertex-transitive and $\alpha(G) = l$. It implies that

$$\chi_f(G) = \frac{|V(G)|}{\alpha(G)} = 4 + \frac{2}{l} \leq 5 - \frac{1}{3}.$$

If $G \neq C_{2l+1} \boxtimes K_2$, then by Lemma 5, we can find an independent set meeting every copy of K_4 and C_8^2 ; we extend it as a maximal independent set I . Note that $G - I$ contains no induced subgraph isomorphic C_8^2 . We have $\chi_f(G - I) \leq 4 - f(4)$; it implies $\chi_f(G) \leq 5 - f(4)$. Thus, $f(3) \geq \min\{f(4), 1/3\}$ and we are finished. $\square$

3 The case $\Delta(G) = 4$

To prove $f(4) \geq \frac{2}{67}$, we will use an approach which is similar to those in [3, 6]. It suffices to prove that the minimum counterexample does not exist.

Let G be a graph with the smallest number of vertices and satisfying

1. $\Delta(G) = 4$ and $\omega(G) \leq 3$;
2. $\chi_f(G) > 4 - \frac{2}{67}$;

3. $G \neq C_8^2$.

By the minimality of G , each vertex in G has degree either 4 or 3. To prove Lemma 1, we will show $\chi_f(G) \leq 4 - \frac{2}{67}$, which gives us the desired contradiction.

For a given vertex x in $V(G)$, it is easy to color its neighborhood $\Gamma_G(x)$ using 2 colors. If $d_G(x) = 3$, then we pick a non-edge S from $\Gamma_G(x)$ and color the two vertices in S using color 1. If $d_G(x) = 4$ and $\alpha(\Gamma_G(x)) \geq 3$, then we pick an independent set S in $\Gamma_G(x)$ of size 3 and assign the color 1 to each vertex in S . If $d_G(x) = 4$ and $\alpha(\Gamma_G(x)) = 2$, then we pick two disjoint non-edges S_1 and S_2 from $\Gamma_G(x)$; we assign color 1 to each vertex in S_1 and color 2 to each vertex in S_2 .

The following Lemma shows that G has a key property, which eventually implies that this local coloring scheme works simultaneously for x in a large subset of $V(G)$.

Lemma 6 *For each $x \in V(G)$ with $d_G(x) = 4$ and $\alpha(\Gamma_G(x)) = 2$, there exist two vertex-disjoint non-edges $S_1(x), S_2(x) \subset \Gamma_G(x)$ satisfying the following property. If we contract $S_1(x)$ and $S_2(x)$, then the resulted graph $G/S_1(x)/S_2(x)$ contains neither K_5^- nor G_0 . Here K_5^- is the graph obtained from K_5 by removing one edge and G_0 is a graph defined in Figure 9.*




Figure 9: The graph G_0 .

The proof of this lemma is quite long and we will present its proof in section 4.

For each vertex x in G , we associate a small set of vertices $S(x)$ selected from $\Gamma_G(x)$ as follows. If $d_G(x) = 3$, then let $S(x)$ be the endpoints of a non-edge in $\Gamma_G(x)$ and label the vertices in $S(x)$ as 1; if $d_G(x) = 4$ and $\alpha(\Gamma_G(x)) \geq 3$, then let $S(x)$ be any independent set of size 3 in $\Gamma_G(x)$ and label all vertices in $S(x)$ as 1; if $d_G(x) = 4$ and $\alpha(\Gamma_G(x)) = 2$, then let $S(x) = S_1(x) \cup S_2(x)$, where $S_1(x)$ and $S_2(x)$ are guaranteed by Lemma 6; we label the vertices in $S_1(x)$ as 1 and the vertices in $S_2(x)$ as 2. For any $x \in V(G)$, we have $|S(x)| = 2, 3$, or 4.

The following definitions depend on the choice of $S(*)$, which is assumed to be fixed through this section. For $v \in G$ and $j \in \{1, 2, 3\}$, we define

$$N_G^j(v) = \{u \mid \text{there is a path } vv_0 \dots v_{j-2}u \text{ in } G \text{ of length } j \text{ such that } v_0 \in S(v) \text{ and } v_{j-2} \in S(u)\}.$$

For $j = 4$, $v \in N_G^4(u)$ if $d_G(u) = 4$, $\alpha(\Gamma_G(u)) = 2$, u and v are connected as shown in Figure 10; otherwise $N_G^4(u) = \emptyset$. In Figure 10, w is connected to one of the two vertices in $S_2(u)$. Similarly, in Figure 11 and 11, a vertex is connected to a group of vertices means it is connected to any vertex in this group.

For $j = 5$, $v \in N_G^5(u)$ if $d_G(w) = 4$, $\alpha(\Gamma_G(w)) = 2$ for $w \in \{u, v\}$ and u and v are connected as shown in Figure 11; otherwise $N_G^5(u) = \emptyset$.

For $j = 7$, $v \in N_G^7(u)$ if $d_G(w) = 4$, $\alpha(\Gamma_G(w)) = 2$ for $w \in \{u, v\}$ and u and v are connected as shown in Figure 12; otherwise $N_G^7(u) = \emptyset$.




Figure 10: 4-th neighborhood. Figure 11: 5-th neighborhood. Figure 12: 7-th neighborhood.

Note that for $j \in \{1, 2, 3, 5, 7\}$, $v \in N_G^j(u)$ if and only if $u \in N_G^j(v)$; but this does not hold for $j = 4$. We have the following lemma.

Lemma 7 *For $u \in V(G)$ such that $d_G(u) = 4$ and $\alpha(\Gamma_G(u)) = 2$, we have $|N_G^1(u) \cup N_G^2(u) \cup N_G^3(u) \cup N_G^4(u) \cup N_G^5(u) \cup N_G^7(u)| \leq 96$.*

Proof: It is clear that $|N_G^1(u) \cup N_G^2(u) \cup N_G^3(u)| \leq 4 + 8 + 8 \times 3 = 36$. We next estimate $|N_G^4(u)|$. In Figure 10, observe that w is connected to one vertices of $S_2(u)$ and $w \notin \Gamma_G(u)$. For a fixed u , there are at most four choices for w , at most three choices for z , and at most three choices for v . Therefore, we have $|N_G^4(u)| \leq 4 \times 3 \times 3 = 36$.

Let us estimate $|N_G^5(u)|$. In Figure 11, for a fixed u , we have four choices for w and two choices for z . Fix a z . Assume $\Gamma_G(z) \setminus \{w\} = \{a, b, c\}$. Let $T_1 = \{a, b\}$, $T_2 = \{b, c\}$, and $T_3 = \{a, c\}$. We have the following claim.

Claim There are at most three $v \in N_G^5(u)$ such that for each v we have $\Gamma_G(z) \cap \Gamma_G(v) = T_i$ for some $1 \leq i \leq 3$ as shown in Figure 11.

Proof of the claim: For each $1 \leq i \leq 3$, there are at most three $v \in N_G^5(u)$ such that $\Gamma_G(z) \cap \Gamma_G(v) = T_i$ as shown in Figure 11 since each vertices in T_i has at most three neighbors other than z . If the claim is false, then there is $1 \leq i \neq j \leq 3$ such that $\Gamma_G(z) \cap \Gamma_G(v_i) = T_i$ and $\Gamma_G(z) \cap \Gamma_G(v'_i) = T_i$ for some $v_i, v'_i \in N_G^5(u)$, and $\Gamma_G(z) \cap \Gamma_G(v_j) = T_j$ for some $v_j \in N_G^5(u)$, where v_i, v'_i, v_j are distinct. Without loss of generality, we assume $\Gamma_G(z) \cap \Gamma_G(v_1) = \Gamma_G(z) \cap \Gamma_G(v'_1) = T_1$ for $v_1, v'_1 \in N_G^5(u)$, and $\Gamma_G(z) \cap \Gamma_G(v_2) = T_2$ for some $v_2 \in N_G^5(u)$, see Figure 13. Observe that $\Gamma_G(b) = \{v_1, v'_1, v_2, z\}$. Since $\Gamma_G(z) \cap \Gamma_G(v_1) = T_1$ as shown in Figure 11, a and one of b 's neighbors form $S_i(v_1)$ for some $i \in \{1, 2\}$; we assume it is $S_1(v_1)$. Note $\{z, v_1, v'_1\} \subset \Gamma_G(a)$. Thus $S_1(v_1) = \{a, v_2\}$ and $v_2 \in \Gamma_G(v_1)$. Similarly, we can show $S_1(v'_1) = \{a, v_2\}$ and $v_2 \in \Gamma_G(v'_1)$. Now, observe that $\Gamma_G(v_2) = \{v_1, v'_1, b, c\}$. Since $\Gamma_G(z) \cap \Gamma_G(v_2) = T_2$ as shown in Figure 11, b and one of neighbors of v_2 form $S_i(v_2)$ for some $i \in \{1, 2\}$; we assume $i = 1$. Because $\{v_1, v'_1\} \subset \Gamma_G(b)$, then $S_1(v_2) = \{b, c\}$. However, b and c are not in the same independent set in the definition of $N_G^5(u)$, see Figure 11. This is a contradiction and this case can not happen. The claim follows.

Therefore, $|N_G^5(u)| \leq 4 \times 2 \times 3 = 24$.

In Figure 12, for a fixed u , we have two choices for the edge e , one choice for w , two choices for z , and three choices for the edge f . Fix a z . By considering the degrees of the endpoints of f , there is at most one f and at most one $v \in N_G^7(u)$ such that $|\Gamma_G(f) \cap \Gamma_G(v)| = 4$ as shown in Figure 12. Therefore, we have $|N_G^7(u)| \leq 2 \times 2 \times 1 = 4$.




Figure 13: The picture for the claim.

Last, we estimate $|N_G^5(u) \cup N_G^7(u)|$. If there is some $v \in N_G^7(u)$, then we observe that there are at most five z 's (see Figure 11). We get the number of $v \in N_G^5(u)$ is at most $5 \times 3 = 15$. In this case, we have

$$|N_G^5(u) \cup N_G^7(u)| \leq 4 + 15 < 24.$$

If $N_G^7(u) = \emptyset$, then also we have

$$|N_G^5(u) \cup N_G^7(u)| \leq 24.$$

Therefore

$$|N_G^1(u) \cup N_G^2(u) \cup N_G^3(u) \cup N_G^4(u) \cup N_G^5(u) \cup N_G^7(u)| \leq 36 + 36 + 24 = 96.$$

□

Based on the graph G , we define an auxiliary graph G^* on vertex set $V(G)$. The edge set is following: $uv \in E(G^*)$ if either $u \in N_G^1(v) \cup N_G^2(u) \cup N_G^3(u) \cup N_G^5(u) \cup N_G^7(u)$, or $v \in N_G^4(u)$, or $u \in N_G^4(v)$. We have the following lemma.

Lemma 8 *The graph G^* is 133-colorable.*

Proof: Let σ be an increasing order of $V(G^*)$ satisfying the following conditions.

- 1: For u and v such that $d_G(u) = 3$ and $d_G(v) = 4$, we have $\sigma(u) < \sigma(v)$.
- 2: For u and v such that $d_G(u) = d_G(v) = 4$, $\alpha(\Gamma_G(u)) \geq 3$, and $\alpha(\Gamma_G(v)) = 2$, we have $\sigma(u) < \sigma(v)$.

We will color $V(G^*)$ according to the order σ . For each v , we have the following estimate on the number of colors forbidden to use for v .

- 1: For v such that $d_G(v) = 3$, the number of colors forbidden to use for v is at most $|N_G^1(v) \cup N_G^2(v) \cup N_G^3(v)| \leq 4 + 8 + 24 = 36$.
- 2: For v such that $d_G(v) = 4$ and $\alpha(\Gamma_G(v)) \geq 3$, the number of colors forbidden to use for v is at most $|N_G^1(v) \cup N_G^2(v) \cup N_G^3(v)| \leq 4 + 8 + 24 = 36$.
- 3: For v such that $d_G(v) = 4$ and $\alpha(\Gamma_G(v)) = 2$, the number of colors forbidden to use for v is at most $|N_G^1(v) \cup N_G^2(v) \cup N_G^3(v) \cup N_G^4(v) \cup N_G^5(v) \cup N_G^7(v)| + |N_G^4(v)| \leq 96 + 36 = 132$ by Lemma 7.

Therefore, The greedy algorithm shows G^* is 133-colorable. □

Let X be a coloring class of G^* . We define a new graph $G(X)$ by the following process.

1. For each $x \in X$, if $|S(x)| = 2$ or $|S(x)| = 3$, then we contract $S(x)$ as a single vertex, delete the vertices in $\Gamma_G(v) \setminus S(v)$, and keep labeling the new vertex as 1; if $|S(x)| = 4$, i.e., $S(x) = S_1(x) \cup S_2(x)$, then we contract $S_1(x)$ and $S_2(x)$ as single vertices and keep their labels. After that, we delete X . Let H be the resulted graph.
2. Note that $\Gamma_H(x) \cap \Gamma_H(y) = \emptyset$ and there is no edge from $\Gamma_H(x)$ to $\Gamma_H(y)$ for any $x, y \in X$ as X is a coloring class.
3. We identify all vertices with label i as a single vertex w_i for $i \in \{1, 2\}$. Let $G(X)$ be the resulted graph.

We have the following lemma on the chromatic number of $G(X)$.

Lemma 9 *The graph $G(X)$ is 4-colorable for each coloring class.*

We postpone the proof of this lemma until the end of this section and prove the Lemma 1 first.

Proof of Lemma 1: By Lemma 8, there is a proper 133-coloring of G^* . We assume $V(G^*) = V(G) = \cup_{i=1}^{133} X_i$, where X_i is the i -th coloring class.

For each $i \in \{1, \dots, 133\}$, Lemma 9 shows $G(X_i)$ is 4-colorable; let $c_i : V(G(X_i)) \rightarrow T_i$ be a proper 4-coloring of the graph $G(X_i)$. Here $T_1, T_2, \dots, T_{133}$ are pairwise disjoint; each of them consists of 4 colors. For $i \in \{1, \dots, 133\}$, the 4-coloring c_i can be viewed as a 4-coloring of $G \setminus X_i$ since each vertex with label j receives the color $c_i(w_i)$ for $j = 1, 2$ and each removed vertex has at most three neighbors in $G \setminus X_i$.

Now we reuse the notation c_i to denote this 4-coloring of $G \setminus X_i$. For each $v \in X_i$, we have $|\cup_{u \in \Gamma_G(v)} c_i(u)| \leq 2$. We can assign two unused colors, denoted by the set $Y(v)$, to v . We define $f_i : V(G) \rightarrow \mathcal{P}(T_i)$ (the power set of T_i) satisfying

$$f_i(v) = \begin{cases} \{c_i(v)\} & \text{if } v \in V(G) \setminus X_i, \\ Y(v) & \text{if } v \in X_i. \end{cases}$$

Observe that each vertex in X_i receives two colors from f_i and every other vertex receives one color. Let $\sigma : V(G) \rightarrow \mathcal{P}(\cup_{i=1}^k T_i)$ be a mapping such that $\sigma(v) = \cup_{i=1}^m f_i(v)$. It is easy to verify σ is a 134-fold coloring of G such that each color is drawn from a palette of 532 colors; namely we have

$$\chi_f(G) \leq \frac{532}{134} = 4 - \frac{2}{67}.$$

The proof of Lemma 1 is finished. □

Before we prove Lemma 9, we need the following definitions.

A *block* of a graph is a maximal 2-connected induced subgraph. A *Gallai tree* is a connected graph in which all blocks are either complete graphs or odd cycles. A *Gallai forest* is a graph all of whose components are Gallai trees. A *k-Gallai tree (forest)* is a Gallai tree (forest) such that the degree of all vertices are at most $k - 1$. A *k-critical* graph is a graph G whose chromatic number is k and deleting any vertex can decrease the chromatic number. Gallai showed the following Lemma.

Lemma 10 [2] *If G is a k -critical graph, then the subgraph of G induced on the vertices of degree $k - 1$ is a k -Gallai forest.*

Proof of Lemma 9: We use proof by contradiction. Suppose that $G(X)$ is not 4-colorable. The only possible vertices in $G(X)$ with degree greater than 4 are the vertices w_1 and w_2 , which are obtained by contracting the vertices with label 1 and 2 in the intermediate graph H . The simple greedy algorithm shows that $G(X)$ is always 5-colorable. Let $G'(X)$ be a

5-critical subgraph of $G(X)$. Applying Lemma 10 to $G'(X)$, the subgraph of $G'(X)$ induced on the vertices of degree 4 is a 5-Gallai forest F . The vertex set of F may contain w_1 or w_2 . Delete w_1 and w_2 from F if F contains one of them. Let F' be the resulted Gallai forest. (Any induced subgraph of a Gallai forest is still a Gallai forest.) The Gallai forest F' is not empty. Let T be a connected component of F' and B be a leaf block of T . The block B is either a clique or an odd cycle from the definition of a Gallai tree.

Let v be a vertex in B . As v has at most two neighbors (w_1 and w_2) outside F' in $G(X)$, we have $d_{F'}(v) \geq 2$. If v is not in other blocks of F' , then we have $d_B(v) \geq 2$. It follows that $|B| \geq 3$. Since B is a subgraph of G and G is K_4 -free, the block B is an odd cycle.

Let $v_1 v_2$ be an edge in B such that v_1 and v_2 are not in other blocks. The degree requirement implies $v_i w_j$ are edges in $G(X)$ for all $i, j \in \{1, 2\}$. For $i = 1, 2$, there are vertices $x_i, y_i \in X$ satisfying $S(x_i) \cap \Gamma_G(v_i) \neq \emptyset$ and $S(y_i) \cap \Gamma_G(v_i) \neq \emptyset$; moreover either $|S(x_i)| = 4$ or $|S(y_i)| = 4$ since one of its neighborhood has label 2. Without loss of generality, we assume $|S(x_i)| = 4$ for $i \in \{1, 2\}$. If $x_i \neq y_i$, then $y_i \in N_G^4(x_i)$, i.e., $y_i \in \Gamma_{G^*}(x_i)$; this contradicts X being a coloring class. Thus we have $x_i = y_i$ and $|S(x_i)| = 4$ for $i \in \{1, 2\}$. For $\{i, j\} = \{1, 2\}$, if $x_i \neq y_j$, then $y_i \in N_G^5(x_i)$, i.e., $y_i \in \Gamma_{G^*}(x_i)$; this is a contradiction of X being a coloring class. Thus we have

$$x_1 = x_2 = y_1 = y_2.$$

Let x denote this common vertex above. Then $d_G(x) = 4$ and $\alpha(\Gamma_G(x)) = 2$.

Let v_0 be the only vertex in B shared by other blocks. Since $B - v_0$ is connected, the argument above shows there is a common x for all edges in $B - v_0$. If $\Gamma_{G(X)}(v_0) \cap \{w_1, w_2\} \neq \emptyset$, then there is some vertex $x_0 \in X$ such that $S(x_0) \cup \Gamma_G(v_0) \neq \emptyset$. By the similar argument, we also have $x_0 = x$.

Therefore, x depends only on B . In the sense that for any $y \in X$ and any $v \in B$, if $S(y) \cap \Gamma_G(v) \neq \emptyset$, then $y = x$.

The block B is an odd cycle as we mentioned above. Suppose $|B| = 2r + 1$. Let $v_0, v_1, \dots, v_{2r}$ be the vertices of B in cyclic order and v_0 be the only vertex which may be shared by other block.

Let $x \in X$ be the vertex determined by B . Recall $d_G(x) = 4$ and $\alpha(\Gamma_G(x)) = 2$. Each vertex in $\Gamma(x)$ can have at most 2 edges to B . We get

$$4r \leq |E(B, \Gamma(x))| \leq 8. \quad (4)$$

We have $r \leq 2$. The block B is either a C_5 or a K_3 . We claim both $v_0 w_1$ and $v_0 w_2$ are non-edges of $G(X)$.

If $B = C_5$, then inequality (4) implies that v_0 has no neighbor in $\Gamma(x)$ and the claim holds. If $B = K_3$, then the claim also holds; otherwise $B \cup \{S_1(x), S_2(x)\}$ forms a K_5^- in $G/S_1(x)/S_2(x)$, which is a contradiction to Lemma 6.

Let u_1 and u_2 be the two neighbors of v_0 in other blocks of F' . If u_1 and u_2 are in the same block, then this block is an odd cycle; otherwise, $v_0 u_1$ and $v_0 u_2$ are in two different blocks.

The union of non-leaf blocks of T is a Gallai-tree, denoted by T' . The argument above shows every leaf block of T' must be an odd cycle. Let C be such a leaf block of T' . Now C is an odd cycle, and C is connected to $|C| - 1$ leaf blocks of T . Let B and B' be two leaf blocks of T such that $B \cap C$ is adjacent to $B' \cap C$. Without loss of generality, we may assume B is the one we considered before. By the same argument, B' is an odd cycle of size $2r' + 1$ with $r' \in \{1, 2\}$. Let $v'_0, v'_1, \dots, v'_{2r'}$ be the vertices of B' and v'_0 be the only vertex in $B' \cap C$. For i in $\{1, 2, \dots, 2r'\}$ and j in $\{1, 2\}$, $v'_i w_j$ are edges in $G'(X)$. Similarly, there exists a vertex $x' \in X$ with $d_G(x') = 4$ and $\alpha(\Gamma_G(x')) = 2$ such that $|E(v_i, S_1(x'))| \geq 1$ and $|E(v_i, S_2(x'))| \geq 1$. We must have $x = x'$; otherwise $x' \in N_G^7(x)$, i.e., $x' \in \Gamma_{G^*}(x)$, and this

contradicts the fact that X is a coloring class in D . Now we have $|E(\Gamma(x), B)| \geq 4r$ and $|E(\Gamma(x), B')| \geq 4r'$. By counting the degrees of vertices in $\Gamma(x)$ in G , we have

$$4r + 4r' + 4 + 4 \leq 16.$$

We get $r = r' = 1$. Both B and B' are K_3 's. In this case, $G/S_1(x)/S_2(x)$ contains the graph G_0 , see figure 9. This contradicts Lemma 6.

We can find the desired contradiction; so the lemma follows. $\square$

4 Proof of Lemma 6

In this section, we will prove Lemma 6. We first review a Lemma in [6].

Lemma 11 *Let G be a graph. Suppose that G_1 and G_2 are two subgraphs such that $G_1 \cup G_2 = G$ and $V(G_1) \cap V(G_2) = \{u, v\}$.*

1. *If uv is an edge of G , then we have*

$$\chi_f(G) = \max\{\chi_f(G_1), \chi_f(G_2)\}.$$

2. *If uv is not an edge of G , then we have*

$$\chi_f(G) \leq \max\{\chi_f(G_1), \chi_f(G_2 + uv), \chi_f(G_2/uv)\},$$

where $G_2 + uv$ is the graph obtained from G_2 by adding edge uv and G_2/uv is the graph obtained from G_2 by contracting $\{u, v\}$.

Proof of Lemma 6: Recall that G is a connected K_4 -free graph with minimum number of vertices such that $G \neq C_8^2$ and $\chi_f(G) > 4 - \frac{2}{67}$. Note that G is 2-connected. We will prove it by contradiction.

Suppose Lemma 6 fails for some vertex x in G . Observe $\Gamma_G(x)$ is one of the graphs in Figure 14. Here we assume $\Gamma_G(x) = \{a, b, c, d\}$. Through the proof of the lemma, let S_1 and S_2 be two vertex-disjoint independent sets in $\Gamma_G(x)$, H be a triangle in $V(G) \setminus (\{x\} \cup \Gamma_G(x))$, then say (S_1, S_2, H) is a bad triple if $\{\underline{S}_1, \underline{S}_2, H\}$ contains a K_5^- in $G/S_1/S_2$.




Figure 14: Three possible cases of $\Gamma_G(x)$.

If $\Gamma_G(x) = P_4$, then $\{a, d\}$ and $\{b, c\}$ is the only pair of disjoint non-edges. There is a triangle H with $V(H) = \{y, z, w\}$ such that $(\{a, d\}, \{b, c\}, H)$ is a bad triple. Note that $|E(\{a, b, c, d\}, \{y, z, w\})| = 5$ or 6 . By an exhaustive search, the induced subgraph of G on $\{x, a, b, c, d, y, z, w\}$ is one of the following six graphs (see Figure 15).




Figure 15: If $\Gamma_G(x) = P_4$, then there are six possible induced subgraphs.

If $\Gamma_G(x) = 2e$, then $(\{a, c\}, \{b, d\})$ and $(\{a, d\}, \{b, c\})$ are two pairs of disjoint non-edges. By considering the degrees of vertices in $\Gamma_G(x)$, there is only one triangle H with $V(H) = \{y, z, w\}$ such that $(\{a, c\}, \{b, d\}, H)$ and $(\{a, d\}, \{b, c\}, H)$ are two bad triples. By an exhaustive search, the induced subgraph of G on $\{x, a, b, c, d, y, z, w\}$ is one of the following three graphs (see Figure 16).




Figure 16: If $\Gamma_G(x) = 2e$, then there are three possible induced subgraphs.

It suffices to show that G cannot contain H_i for $1 \leq i \leq 9$. Since all vertices in H_1 (and H_2) have degree 4, H_1 (and H_2) is the entire graph G . Observe that H_1 is isomorphic to C_8^2 and H_2 is 11:3-colorable (see Figure 17). Contradiction!

In H_7 , the vertex d is the only vertex with degree less than 4. If H_7 is not the entire graph G , then d is a cut vertex of G . This contradicts the fact that G is 2-connected. Thus $G = H_7$. The graph H_7 is 11:3-colorable as shown by Figure 17. Contradiction!

Now we consider the case H_3 . Note $H_3 + bz$ is the graph H_2 . We have $\chi_f(H_3) \leq \chi_f(H_2) \leq 11/3$. The graph H_3 must be a proper induced subgraph of G , and the pair $\{b, z\}$ is a vertex cut of G . Let G' be the induced subgraph of G by deleting all vertices in H_3 but b, z . We




Figure 17: The graph H_2 and H_7 are 11:3-colorable.

apply Lemma 11 to $G + bz$ with $G_1 = H_3 + bz = H_2$ and $G_2 = G' + bz$. We have

$$\chi_f(G + bz) \leq \max\{\chi_f(H_2), \chi_f(G' + bz)\}.$$

Note $\chi_f(H_2) \leq 11/3$ and $11/3 < \chi_f(G) \leq \chi_f(G + bz)$. We have $\chi_f(G) \leq \chi_f(G' + bz)$. Both b and z have at most 2 neighbors in $G' + bz$. Thus $G' + bz$ is K_4 -free; $G' + bz \neq C_8^2$ and has fewer vertices than G . This contradicts to the minimality of G .

Note $H_5 + cy = H_2$. The case H_5 is similar to the case H_3 .

Note that H_4 , H_6 , and H_8 are isomorphic to each other. It suffices to show G does not contain H_4 . Suppose that H_4 is a proper induced subgraph of G . Let G_1 be the induced subgraph of G by deleting all vertices in H_4 . Note C_8^2 is not a proper subgraph of any graph in $\mathcal{G}_4$. We have $G_1 \neq C_8^2$. Note that c and z have degree 3 while other vertices in H_4 have degree 4. Since G is 2-connected, c has a unique neighbor, denoted by u , in $V(G_1)$. Similarly, z has a unique neighbor, denoted by v , in $V(G_1)$. Observe that the pair $\{u, v\}$ forms a vertex cut of G . Let G_2 be the induced graph of G on $V(H_4) \cup \{u, v\}$. Applying Lemma 11 to G with G_1 and G_2 , we have

$$\chi_f(G) \leq \max\{\chi_f(G_1), \chi_f(G_2 + uv), \chi_f(G_2/uv)\}.$$

Figure 18 shows $\chi_f(G_2 + uv)$ and $\chi_f(G_2/uv)$ are at most $11/3$.




Figure 18: Case H_4 : both graph $G_2 + uv$ and G_2/uv are 11:3-colorable.

Since $\chi_f(G) > 11/3$, we have $\chi_f(G) \leq \chi_f(G_1)$. Now G_1 is K_4 -free and has maximum degree at most 4; G_1 has fewer vertices than G . This contradicts the minimality of G .

If $G = H_4$, then $\chi_f(H_4) \leq 11/3$, since H_4 is a subgraph of $G_2 + uv$ in Figure 18.

Now we consider the last case H_9 . First, we contract b, c, z into a fat vertex denoted by $\underline{bcz}$. We write G/bcz for the graph after this contraction. Observe that $\{\underline{bcz}, d\}$ is a vertex-cut of G/bcz . Let G_4 and G'_4 be two connected subgraphs of G/bcz such that $G_4 \cup G'_4 = G/bcz$, $G_4 \cap G'_4 = \{\underline{bcz}, d\}$, and $\{u, v\} \subset G'_4$. Note that G_4 is 11:3 colorable, see Figure 19. Now by Lemma 11, we have

$$\chi_f(G/bcz) \leq \max\{\chi_f(G_4), \chi_f(G'_4)\}.$$

As $\{b, c, z\}$ is an independent set, each $a:b$ -coloring of G/bcz gives an $a:b$ -coloring of G , that is $\chi_f(G/bcz) \geq \chi_f(G) > 11/3$. The graph G_4 is 11:3-colorable; see Figure 19. Thus we have $\chi_f(G'_4) \geq \chi_f(G/bcz) \geq \chi_f(G)$. It is easy to check that G'_4 has maximum degree 4, K_4 -free, and it is not C_8^2 . Hence G'_4 must contain a K_4 . Otherwise, it contradicts the minimality of G .

Second, we contract $\{b, d, z\}$ into a fat vertex $\underline{bdz}$ and denote the graph by G/bcz . Let G_5 and G'_5 be two connected subgraphs of G/bdz such that $G_5 \cup G'_5 = G/bdz$, $G_5 \cap G'_5 = \{\underline{bdz}, c\}$, and $\{u, v\} \subset G'_5$. Note that G_5 is 11:3-colorable; see Figure 19. By a similar argument, G'_5 must contain a K_4 .




Figure 19: Case H_9 : the graphs G_4 , G_5 , and G_6 are 11:3-colorable.

The remaining case is that both G'_4 and G'_5 have a K_4 when we contract b and z . Since the origin graph G is K_4 -free, the K_4 in G'_4 (and in G'_5) must contain the fat vertex $\underline{bcz}$ (or $\underline{bdz}$), respectively. Note that each of the four vertices b, c, d, z has at most one edge leaving H_9 . There must be a triangle uvp in G and these four outward edges are connected to some element of $\{u, v, p\}$. The graph G/bz must contain the subgraph G_6 as drawn in Figure 19.

Note that $\{u, v\}$ is a vertex-cut in G/bz . Let G_6 and G'_6 be two connected subgraphs of G/bz , which satisfy $G_6 \cup G'_6 = G$, $G_6 \cap G'_6 = \{u, v\}$, and $\underline{bz} \in G_6$. By Lemma 11, we have

$$\chi_f(G/bz) \leq \max\{\chi_f(G_6), \chi_f(G'_6)\}.$$

Note that G_6 is 11:3-colorable; see Figure 19. We also have $\chi_f(G/bz) \geq \chi_f(G) > \frac{11}{3}$. We obtain $\chi_f(G'_6) \geq \chi_f(G/bz) \geq \chi_f(G)$. Observe that G'_6 is a subgraph of G . We arrive at a contradiction of the minimality of G .

If $\Gamma_G(x) = C_4$, then the only possible choice for the two independent sets are $\{a, c\}$ and $\{b, d\}$. If there is some triangle H such that $(\{a, c\}, \{b, d\}, H)$ is a bad triple, then we have

$$|E(\Gamma_G(x), H)| \geq 5.$$

However, $|E(\Gamma_G(x), H)| \leq 4$. This is a contradiction. Thus the lemma follows in this case.

We can select two vertex disjoint non-edges S_1 and S_2 such that the graph $G/S_1/S_2$ contains no K_5^- . For these particular S_1 and S_2 , if $G/S_1/S_2$ contains no G_0 , then Lemma 6 holds.

Without loss of generality, we assume that $G/S_1/S_2$ does contain G_0 . Let $s_i = \underline{S_i}$ for $i = 1, 2$. Observe that both s_1 and s_2 have four neighbors u, v, p, q other than x in $\bar{G}_0$. It follows that

$$|E(S_1 \cup S_2, \{u, v, p, q\})| \geq 8.$$

On the one hand, we have

$$\begin{aligned}
|E(G|_{S_1 \cup S_2})| &= \frac{1}{2} \left(\sum_{v \in S_1 \cup S_2} d(v) - |E(S_1 \cup S_2, \{u, v, p, q\})| - 4 \right) \\
&\leq \frac{1}{2}(16 - 8 - 4) \\
&= 2.
\end{aligned}$$

On the other hand, $\alpha(\Gamma(x)) = 2$ implies $G|_{S_1 \cup S_2}$ contains at least two edges. Thus, we have $\Gamma_G(x) = 2e$. Label the vertices in $\Gamma_G(x)$ by a, b, c, d as in Figure 14. We assume ab and cd are edges while ac, bd, ad, bc are non-edges. Observe that each vertex in $\{u, v, p, q\}$ has exactly two neighbors in $\{a, b, c, d\}$.

If one vertex, say u , has two neighbors forming a non-edge, say ac , then we can choose $S'_1 = \{a, c\}$ and $S'_2 = \{b, d\}$. It is easy to check that $G/S'_1/S'_2$ contains neither G_0 nor K_5^- . We are done in this case.

In the remaining case, we can assume that for each vertex y in $\{u, v, p, q\}$, the neighbors of y in $\{a, b, c, d\}$ always form an edge. Up to relabeling vertices, there is only one arrangement for edges between $\{u, v, p, q\}$ and $\{a, b, c, d\}$; see the graph H_{10} defined in Figure 20. The




Figure 20: H_{10} and an 11:3-coloring of H_{10} .

graph H_{10} is 11:3-colorable as shown in Figure 20. Since $\chi_f(G) > 11/3$, H_{10} is a proper subgraph of G . Note in H_{10} , every vertices except w and r has degree 4; both w and r have degree 3. Thus, $\{w, r\}$ is a vertex cut of G . Let $G_1 = H_{10}$ and G_2 be the subgraph of G by deleting vertices in $\{x, a, b, c, d, p, q, u, v\}$. Applying Lemma 11 with G_1 and G_2 defined above, we have

$$\chi_f(G) \leq \max\{\chi_f(G_1), \chi_f(G_2)\}.$$

Since $\chi_f(G) > 11/3$ and $\chi_f(G_1) \leq 11/3$ (see Figure 20), we must have $\chi_f(G_2) \geq \chi_f(G)$. Note that G_2 has fewer number of vertices than G . This contradicts the minimality of G . Therefore, the lemma follows. $\square$

References

- [1] O. Borodin and A. Kostochka, On an upper bound on a graph's chromatic number, depending on the graph's degree and density, *J. Comb. Th. B.*, **23** (1977), 247-250.
- [2] T Gallai, Kritische graphen I, *Magyar Tud. Akad. Mat. Kutat ó Int. Közl*, **8** (1963), 165-192.

- [3] H. Hatami and X. Zhu, The fractional chromatic number of graphs of maximum degree at most three, *SIAM. J. Discrete Math.*, **24** (2009), 1762-1775.
- [4] C. C. Heckman and R. Thomas, A new proof of the independence ratio of triangle-free cubic graphs, *Discrete Math.*, **233** (2001), 233-237.
- [5] Andrew D. King, Hitting all maximum cliques with a stable set using lopsided independent transversals, *J. of Graph Theory*, **67**-4 (2011), 300-305.
- [6] L. Lu and X. Peng, The fractional chromatic number of triangle-free graphs with $\Delta \leq 3$, *submitted*.
- [7] M. Molloy and B. Reed, *Graph colouring and the probabilistic method*, volume 23 of Algorithms and Combinatorics, Springer-Verlag, Berlin, 2002.
- [8] B. Reed, ω , Δ , and χ , *J. of Graph Theory*, **27**-4 (1998), 177-227.
- [9] B. Reed, A strengthening of Brooks' theorem, *J. Comb. Th. B.*, **76** (1999), 136-149.
- [10] E. R. Scheinerman and D. H. Ullman, *Fractional graph theory*. A rational approach to the theory of graphs, Wiley-Intersci. Ser. Discrete Math. Optim, John Wiley & Sons, Inc, New York, 1997.